

ON THE REGULARITY NUMBER OF A FINITE GROUP AND OTHER BASE-RELATED INVARIANTS

MARINA ANAGNOSTOPOULOU-MERKOURI AND TIMOTHY C. BURNES

ABSTRACT. A k -tuple (H_1, \dots, H_k) of core-free subgroups of a finite group G is said to be regular if G has a regular orbit on the Cartesian product $G/H_1 \times \dots \times G/H_k$. The regularity number of G , denoted $R(G)$, is the smallest positive integer k with the property that every such k -tuple is regular. In this paper, we develop some general methods for studying the regularity of subgroup tuples in arbitrary finite groups, and we determine the precise regularity number of all almost simple groups with an alternating or sporadic socle. For example, we prove that $R(S_n) = n - 1$ and $R(A_n) = n - 2$. We also formulate and investigate natural generalisations of several well-studied problems on base sizes for finite permutation groups, including conjectures due to Cameron, Pyber and Vdovin. For instance, we extend earlier work of Burnes, O'Brien and Wilson by proving that $R(G) \leq 7$ for every almost simple sporadic group, with equality if and only if G is the Mathieu group M_{24} . We also show that every triple of soluble subgroups in an almost simple sporadic group is regular, which generalises recent work of Burnes on base sizes for transitive actions of sporadic groups with soluble point stabilisers.

CONTENTS

1.	Introduction	1
2.	Preliminaries	7
2.1.	Probabilistic methods	7
2.2.	Independent sets	9
2.3.	Almost simple groups	10
2.4.	Computational methods	13
3.	Symmetric and alternating groups	14
3.1.	Primitive subgroups	14
3.2.	The regularity number	19
4.	Sporadic groups	33
4.1.	The regularity number	34
4.2.	Soluble subgroups	42
	References	50

1. INTRODUCTION

Let G be a finite group and let H_1, \dots, H_k be a collection of core-free subgroups of G , allowing repetitions. Consider the natural action of G on the Cartesian product

$$X = G/H_1 \times \dots \times G/H_k$$

and observe that G has a regular orbit on X if and only if

$$\bigcap_{i=1}^k H_i^{g_i} = 1$$

for some elements $g_i \in G$. We introduce the following definition.

Date: May 23, 2024.

Definition. A k -tuple (H_1, \dots, H_k) of core-free subgroups of G is *regular* if G has a regular orbit on X , and *non-regular* otherwise.

For $k = 2$ or 3 , we will refer to regular and non-regular pairs or triples, respectively. Clearly, the regularity of a given tuple is independent of the ordering of the subgroups in the tuple. In addition, (H_1, \dots, H_k) is regular if and only if $(H_1^{g_1}, \dots, H_k^{g_k})$ is regular for some $g_i \in G$, so we are free to replace each H_i by any conjugate. Given some group-theoretic property \mathcal{P} of a subgroup of G , such as maximal, soluble or nilpotent, we will say that (H_1, \dots, H_k) has property \mathcal{P} if every component subgroup in the tuple has this property (we will always assume that the subgroups in any given tuple are core-free). It will also be convenient to say that such a k -tuple is *conjugate* if each H_i is conjugate to H_1 .

Recall that if $G \leq \text{Sym}(\Omega)$ is a permutation group on a set Ω , then a subset of Ω is a *base* if its pointwise stabiliser in G is trivial. So a conjugate tuple (H_1, \dots, H_k) is regular if and only if G has a base of size k with respect to its natural action on G/H_1 . Of course, the latter is equivalent to the bound $k \geq b(G, H_1)$, where $b(G, H_1)$ is the *base size* of G , which is the minimal size of a base for the action of G on G/H_1 . There is a very substantial literature on bases for finite permutation groups stretching all the way back to the nineteenth century, finding an extensive range of applications and connections to other areas of group theory and combinatorics (we refer the reader to the survey articles [4, 37, 39] and [8, Section 5] for further details). In view of this connection, we introduce the following invariants.

Definition. Let G be a finite group.

- (i) The *regularity number* of G , denoted $R(G)$, is the minimal positive integer k such that every k -tuple of core-free subgroups of G is regular.
- (ii) The *base number* of G , denoted $B(G)$, is the minimal positive integer k such that $b(G, H) \leq k$ for every core-free subgroup H of G .

Remark 1. Let us record some immediate remarks on these definitions:

- (a) Clearly, we have $B(G) \leq R(G)$. It turns out that there are examples where this inequality is strict. For instance, if G is the sporadic simple group M_{11} , then the main theorem of [16] gives $B(G) = 4$, but we find that $R(G) = 5$. Indeed, G has maximal subgroups $H = M_{10} = A_6.2$ and $K = L_2(11)$, and one can check that the 4-tuple (H, H, K, K) is non-regular. In fact, we will show that there are infinitely many finite simple groups with $R(G) \geq B(G) + 1$ (see Proposition 2.12). Moreover, in Proposition 2.13 we prove that if $G = \text{GL}_n(2)$ with $n \geq 5$, then $B(G) = n$ and $R(G) \geq 2(n - 1)$, so the difference $R(G) - B(G)$ can be arbitrarily large.
- (b) In [19], Cameron introduces several base-related invariants of a finite group G . Let us define $b_2(G)$ to be the maximum, over all faithful permutation representations of G , of the maximal size of a minimal base (recall that a base is *minimal* if no proper subset is a base). Since $B(G)$ is the minimal size of a minimal base over all faithful *transitive* permutation representations of G , we have $B(G) \leq b_2(G)$. In [19, Corollary 3.3], Cameron proves that $b_2(G) \leq \sigma(G)$ (also see Proposition 2.3), where $\sigma(G)$ is the maximal size of an *independent subset* of G (that is, $\sigma(G)$ is the maximal size of a subset S of G such that $x \notin \langle S \setminus \{x\} \rangle$ for all $x \in S$). By combining this result with a theorem of Whiston [53] on independent subsets of symmetric groups, it follows that

$$B(G) = b_2(G) = \sigma(G) = n - 1$$

for $G = S_n$ (see Corollary 2.4).

- (c) Two related invariants of a finite group were studied in [11, 12]. The *intersection number* of G is defined to be the minimal number of maximal subgroups of G whose intersection coincides with the Frattini subgroup of G . And for a group with trivial Frattini subgroup, the *base number* was defined in [12] to be the minimal base size

over all faithful primitive permutation representations of G . The main results in [11, 12] include sharp upper bounds on these invariants for almost simple groups.

A wide range of related invariants can be defined in this setting. Indeed, if \mathcal{P} is a group-theoretic property satisfied by certain core-free subgroups of G (such as maximality, solubility, nilpotency, etc.) then we can define $R_{\mathcal{P}}(G)$ and $B_{\mathcal{P}}(G)$ by restricting to tuples comprising subgroups with property \mathcal{P} . The flexibility of this set-up allows us to succinctly describe a number of well-studied problems concerning bases for transitive groups, leading to natural extensions and open problems in the more general regularity setting. Let us highlight three main examples:

Vdovin's conjecture. Let G be a finite group with trivial soluble radical and let $B_{\text{sol}}(G)$ be the maximal base size $b(G, H)$ over all core-free soluble subgroups H of G . Then a conjecture of Vdovin (see [41, Problem 17.41(b)]) asserts that $B_{\text{sol}}(G) \leq 5$, which is a strengthening of an earlier conjecture of Babai, Goodman and Pyber (see Conjecture 6.6 in [2]). The example $G = S_8$ with $H = S_4 \wr S_2$ shows that the bound in Vdovin's conjecture would be best possible (in fact, there are infinitely many examples with $b(G, H) = 5$). Although Vdovin's conjecture remains open, there has been some recent progress towards a positive solution. For example, the main theorem of [9] establishes the bound $B_{\text{sol max}}(G) \leq 5$ with respect to soluble maximal subgroups, and Vdovin [48] has reduced the conjecture to almost simple groups (recall that a finite group G is *almost simple* if there exists a non-abelian simple group T such that $T \triangleleft G \leq \text{Aut}(T)$; here T is the *socle* of G). In the latter setting, the conjecture for groups with an alternating or sporadic socle is resolved in [3, 10], but the general problem for groups of Lie type remains open.

Cameron's conjecture. In a different direction, a highly influential conjecture of Cameron and Kantor [21] from 1993 asserts that there exists an absolute constant c such that $B_{\text{ns}}(G) \leq c$ for every almost simple group G . Here $B_{\text{ns}}(G)$ denotes the maximal base size $b(G, H)$ over all *non-standard* maximal subgroups H of G (roughly speaking, a maximal subgroup H is *standard* if G is either a classical group and H acts reducibly on the natural module, or $G = S_n$ or A_n and H acts intransitively or imprimitively on $\{1, \dots, n\}$, otherwise H is *non-standard*). This conjecture was proved by Liebeck and Shalev [36] (with an undetermined constant) using a powerful probabilistic approach for studying bases. In response, Cameron [20, p.122] conjectured that 7 is the optimal bound, with $B_{\text{ns}}(G) = 7$ if and only if $G = M_{24}$. Cameron's conjecture was proved in a series of papers by Burness et al. [7, 14, 15, 16].

Pyber's conjecture. Another intensively studied conjecture on base sizes was proposed by Pyber in the early 1990s [46, p.207]. Before stating the conjecture, first observe that if G is a finite transitive permutation group of degree n with point stabiliser H , then

$$b(G, H) \geq \log_n |G|.$$

Pyber's conjecture asserts that all primitive groups admit small bases in the sense that there exists an absolute constant c such that $b(G, H) \leq c \log_n |G|$ for every finite primitive group G of degree n . Building on earlier work by several authors (see [17, 38, 47], for example), the proof of Pyber's conjecture was completed by Duyan et al. in [27]. This was extended in [32], where the main theorem gives the explicit bound

$$b(G, H) \leq 2 \log_n |G| + 24.$$

Turning to the base number, suppose G is a finite group with a core-free maximal subgroup and let $m(G)$ be the minimal index of such a subgroup (in other words, $m(G)$ is the minimal degree of a faithful primitive permutation representation of G). Then

$$B_{\text{max}}(G) \geq \log_{m(G)} |G|,$$

where $B_{\text{max}}(G)$ is the maximal base size of G over all primitive faithful permutation representations (and we can define $R_{\text{max}}(G)$ with respect to tuples of core-free maximal subgroups

of G). And the positive solution to Pyber's conjecture implies that $B_{\max}(G) \leq c \log_{m(G)} |G|$ for some absolute constant c .

Here we propose natural extensions of the above base size conjectures of Pyber, Cameron and Vdovin in the more general regularity setting.

Conjecture 1.

- (i) *There exists an absolute constant c such that*

$$R_{\max}(G) \leq c \log_{m(G)} |G|$$

for every finite group G with a core-free maximal subgroup.

- (ii) *We have $R_{\text{ns}}(G) \leq 7$ for every finite almost simple group G , with equality if and only if $G = M_{24}$.*
- (iii) *We have $R_{\text{sol}}(G) \leq 5$ for every finite group G with trivial soluble radical.*

In this paper, we develop some general techniques for studying the regularity number of a finite group and we take the first steps towards establishing the above conjectures for almost simple groups with socle an alternating or sporadic group. These results will be extended to groups of Lie type in a follow up paper by Anagnostopoulou-Merkouri. We now state our main results, first focussing on symmetric and alternating groups.

Let G be an almost simple group with socle $T = A_n$. We say that a core-free subgroup H of G is *primitive* if $H \cap T$ acts primitively on the set $\{1, \dots, n\}$, and we define $B_{\text{prim}}(G)$ and $R_{\text{prim}}(G)$ accordingly. Similarly, we define $R_{\text{intrans}}(G)$ and $R_{\text{sol max}}(G)$ in terms of tuples of intransitive and soluble maximal subgroups of G , respectively. (Of course, if G does not have a core-free primitive subgroup, then $B_{\text{prim}}(G)$ and $R_{\text{prim}}(G)$ are undefined, and similarly for $R_{\text{sol max}}(G)$. For example, no proper subgroup of A_{34} acts primitively on $\{1, \dots, 34\}$.)

Theorem 1. *Let G be an almost simple group with socle $T = A_n$.*

- (i) *For $G \in \{S_n, A_n\}$ we have $R(G) = R_{\text{intrans}}(G) = B(G) = n - |S_n : G|$.*
- (ii) *Suppose G has a core-free primitive subgroup. Then $R_{\text{prim}}(G) \leq 6$, with equality if and only if $G = A_8$. Moreover, if $n \geq 13$, then $R_{\text{prim}}(G) = 2$.*
- (iii) *Suppose G has a soluble maximal subgroup. Then $R_{\text{sol max}}(G) \leq 5$, with equality if and only if $G = S_8$. Moreover, if $n \geq 17$, then $R_{\text{sol max}}(G) = 2$.*

Remark 2. Let us record some comments on the statement of Theorem 1.

- (a) For $T = A_6$ we note that $R(G) = B(G) = 4$ if $G = \text{PGL}_2(9)$, M_{10} or $A_6.2^2$.
- (b) Suppose $G = S_n$. Clearly, if $H = S_{n-1}$ is intransitive then the $(n-2)$ -tuple (H, \dots, H) is non-regular since the intersection of any $n-2$ conjugates of H will contain a transposition. More generally, we can determine all the maximal intransitive non-regular $(n-2)$ -tuples for S_n (see Proposition 3.21); up to ordering and conjugacy, they are of the form (H, \dots, H, K) , where $H = S_{n-1}$ and $K = S_k \times S_{n-k}$ for some $1 \leq k \leq n/2$. We refer the reader to Remark 3.23 for further comments.
- (c) Notice that the natural actions of S_n and A_n on $\{1, \dots, n\}$ immediately give the lower bounds $B(S_n) \geq n-1$ and $B(A_n) \geq n-2$. So as a corollary to Theorem 1, we deduce that $B(S_n) = n-1$ and $B(A_n) = n-2$. Following Cameron [19], we can also show that $B(S_n) = n-1$ by combining the bound $B(S_n) \leq \sigma(S_n)$ with Whiston's theorem on independent subsets in [53], as discussed in Remark 1(b). Similarly, we get $B(A_n) = n-2$ (see Corollary 2.4).
- (d) We have $m(G) = n$ and thus Theorem 1(i) yields

$$R(G) = n - |S_n : G| < 2 \log_{m(G)} |G|$$

for all $G \in \{S_n, A_n\}$ with $n \geq 5$. In particular, this establishes part (i) of Conjecture 1 for all symmetric and alternating groups.

- (e) The bound in (ii) extends the main theorem of [14] by establishing a stronger form of Cameron's conjecture (see Conjecture 1(ii)) for all almost simple groups with socle an alternating group. We refer the reader to Table 1 in Section 3.1 for a complete list of the groups G with $R_{\text{prim}}(G) = k > 2$, together with an example of a non-regular primitive $(k - 1)$ -tuple. The exact value of $B_{\text{prim}}(G)$ is determined in [14] and we deduce that $R_{\text{prim}}(G) \neq B_{\text{prim}}(G)$ if and only if $G = A_8$.
- (f) Part (iii) extends the main theorem of [9] for groups with socle A_n and it establishes a special case of Conjecture 1(iii). We have computed the exact value of $R_{\text{sol max}}(G)$ in all cases and the groups with $R_{\text{sol max}}(G) = k > 2$ are recorded in Table 2 (see Section 3.1) along with an example of a non-regular $(k - 1)$ -tuple of the form (H, \dots, H) . In particular, we have $R_{\text{sol max}}(G) = B_{\text{sol max}}(G)$ for all $n \geq 5$.

In [3], Baykalov proves that $B_{\text{sol}}(G) \leq 5$ for every almost simple group G with socle A_n and a special case of Conjecture 1(iii) asserts that $R_{\text{sol}}(G) \leq 5$. With the aid of MAGMA [5], we have verified the latter bound for the groups with $n \leq 11$ and we find that $R_{\text{sol}}(G) = 5$ if and only if $G = S_8$ (and up to conjugacy, the only non-regular soluble 4-tuple is (H, H, H, H) with $H = S_4 \wr S_2$). We speculate that the following much stronger conjecture holds in this setting.

Conjecture 2. *We have $R_{\text{sol}}(S_n) = R_{\text{sol}}(A_n) = 2$ for all sufficiently large n .*

As recorded in Table 2, we have $R_{\text{sol max}}(G) = 3$ for $G = S_{16}$ since $b(G, H) = 3$ when $H = S_4 \wr S_4$. More generally, if $G = S_{16+m}$ and $H = (S_4 \wr S_4) \times S_m < G$ with $m \leq 4$, then with the aid of MAGMA [5] we can show that $b(G, H) = 3$. We also note that $b(G, H) = 2$ when $G = S_{21}$ and $H = (S_4 \wr S_4) \times S_4$, which leads us to speculate that $R_{\text{sol}}(S_n) = R_{\text{sol}}(A_n) = 2$ for all $n \geq 21$. In the context of the above conjecture, it is also worth recalling a theorem of Dixon [26], which states that $|H| \leq 24^{(n-1)/3}$ for every soluble subgroup H of $G = S_n$ (with equality if $n = 4^d$ and $H = S_4 \wr S_4 \wr \dots \wr S_4$ is the iterated wreath product of d copies of S_4). In particular, Dixon's bound implies that $|H|^3 < |G|$ for all $n \geq 59$.

For nilpotent subgroups, Zenkov [56] uses the Classification of Finite Simple Groups to prove that $R_{\text{nilp}}(G) \leq 3$ for every finite group G with trivial Fitting subgroup. This bound is best possible (for instance, if $G = S_8$ and H is a Sylow 2-subgroup, then $b(G, H) = 3$) and interest in this problem can be traced all the way back to work of Passman in the 1960s. For example, in [45], Passman proves that if G is a finite p -soluble group and P is a Sylow p -subgroup, then there exist $x, y \in G$ such that $P \cap P^x \cap P^y = O_p(G)$ is the p -core of G .

In [54], Zenkov proves that every nilpotent pair of subgroups of S_n or A_n (for $n \geq 5$) is regular, with the single exception of the symmetric group S_8 . Computations in the low degree groups lead us to propose the following conjecture.

Conjecture 3. *If $G = S_n$ or A_n with $n \geq 13$, then every pair of subgroups (H, K) is regular when H is nilpotent and K is soluble.*

Note that $G = S_{12}$ has a non-regular pair (H, K) , where H is a Sylow 2-subgroup and $K = S_4 \wr S_3$, so the conjectured bound $n \geq 13$ would be best possible.

Now let us state our main result for almost simple sporadic groups, which establishes a strong form of Conjecture 1 in this setting (in part (ii), $c = 1$ or 2).

Theorem 2. *Let G be an almost simple sporadic group with socle T .*

- (i) *The exact values of $B(G)$ and $R(G)$ are recorded in Table 3. In particular, $R(G) \leq 7$, with equality if and only if $G = M_{24}$.*
- (ii) *We have $R_{\text{sol}}(G) \leq 3$, with equality if*

$$G = M_{11}, M_{12}.c, M_{22}.c, M_{23}, M_{24}, J_2.c, \text{HS}.2, \text{Co}_2, \text{Fi}_{22}.c \text{ or } \text{Fi}_{23}.$$

- (iii) Suppose G has a soluble maximal subgroup. Then $R_{\text{sol max}}(G) \leq 3$, with equality if and only if $T = M_{11}, M_{12}, J_2, Fi_{22}$ or Fi_{23} .

Remark 3. Some comments on the statement of Theorem 2 are in order.

- (a) In part (i), the base number of every almost sporadic group G can be read off from the main theorem of [16], which gives $b(G, H)$ for every core-free maximal subgroup H . As an immediate consequence of part (i), we observe that $R(G) \leq B(G) + 1$, with equality if and only if $G = M_{11}$ or M_{12} .
- (b) In Table 3 (see Section 4.1) we list all the *large* non-regular k -tuples with $k = R(G) - 1$, up to conjugacy and ordering (large tuples are defined in Definition 4.1), with the exception of the Baby Monster and the Monster (see Remark 4.11 for comments on the latter two cases). It follows that the only non-regular 6-tuple arises when $G = M_{24}$ and every component is conjugate to $H = M_{23}$.
- (c) A more detailed result on $R_{\text{sol}}(G)$ is given in Proposition 4.12. In particular, we have computed $R_{\text{sol}}(G)$ precisely unless T is one of the following groups:

$$Co_1, HN, J_4, Ly, Th, Fi'_{24}, B, M.$$

In addition, for some of the groups with $R_{\text{sol}}(G) = 3$ we present a non-regular soluble pair in Table 8. The complete list of non-regular soluble maximal pairs is recorded in Table 10 (see Proposition 4.20).

- (d) In part (iii), note that every almost simple sporadic group G has a soluble maximal subgroup unless $G = M_{12}, M_{12.2}, M_{24}$ or HS. For the latter groups, $R_{\text{sol max}}(G)$ is not defined.
- (e) We refer the reader to Remark 4.22 for some comments on the existence of non-regular pairs (H, K) with H nilpotent and K soluble. It is worth noting that we are only aware of examples for the groups $M_{22.2}$ and $J_2.2$.

We conclude this introduction by briefly commenting on the proofs of Theorems 1 and 2, which involve a combination of probabilistic, combinatorial and computational methods.

Let G be a finite group, let $\tau = (H_1, \dots, H_k)$ be a k -tuple of core-free subgroups and let

$$Q(G, \tau) = \frac{|\{(\alpha_1, \dots, \alpha_k) \in \Gamma : \bigcap_{i=1}^k G_{\alpha_i} \neq 1\}|}{|\Gamma|}$$

be the probability that a randomly chosen tuple in $\Gamma = G/H_1 \times \dots \times G/H_k$ is not contained in a regular orbit of G , with respect to the uniform distribution on Γ . Notice that τ is regular if and only if $Q(G, \tau) < 1$.

By adapting earlier work of Liebeck and Shalev [36] on base sizes, we can estimate $Q(G, \tau)$ in terms of fixed point ratios (see Lemma 2.1) and this provides a powerful tool for establishing the regularity of τ . Indeed, this approach is at the heart of our proof of parts (ii) and (iii) in Theorem 1, where we can appeal to earlier work of Guralnick and Magaard [30] and Maróti [40] in order to derive appropriate bounds on the relevant fixed point ratios. It is also a key ingredient in our proof of Theorem 2, especially in our analysis of some of the larger sporadic groups. For example, if $G = M$ is the Monster then we can use the GAP Character Table Library [6] to show that $Q(G, \tau) < 1$ for every maximal triple τ , and by combining this with the main theorem of [16] we deduce that $R(M) = 3$ and $R_{\text{sol}}(M) \leq 3$.

In stark contrast, our proof of part (i) of Theorem 1 for $G = S_n$ or A_n is essentially constructive, and it relies on a highly combinatorial analysis of the stabilisers of subsets and partitions of $\{1, \dots, n\}$. Here one of the key results is Lemma 3.15, which shows that if H_1 and H_2 are the stabilisers in G of uniform partitions, then we can typically find an element $g \in G$ such that $H_1 \cap H_2^g$ fixes any given 2-element subset.

Our proof of Theorem 2 on sporadic groups is entirely computational, working with MAGMA [5] and GAP [28]. For example, if $\tau = (H_1, \dots, H_k)$ is a maximal k -tuple then

we can often use MAGMA to construct G and each H_i as subgroups of some symmetric group S_n (typically where $n = m(G)$ is the minimal degree of a faithful primitive permutation representation of G) and then use random search to identify elements $g_i \in G$ with $\bigcap_i H_i^{g_i} = 1$. On the other hand, if we are seeking to show that τ is non-regular, then in many cases we can directly calculate the orbits of G on $G/H_1 \times \cdots \times G/H_k$ in order to confirm the non-existence of a regular orbit. Of course, these computations are not feasible for some of the larger sporadic groups and they require a different approach. For instance, the probabilistic method outlined above is often a very useful tool in this setting, working closely with the detailed information on sporadic groups available in the GAP Character Table Library [6] and the Web Atlas [49].

Notation. Our notation is all fairly standard. Let G be a finite group and let n be a positive integer. We will write C_n , or just n , for a cyclic group of order n and G^n will denote the direct product of n copies of G . An unspecified extension of G by a group H will be denoted by $G:H$; if the extension splits then we may write $G:H$. We adopt the standard notation for simple groups of Lie type from [34].

Organisation. We begin in Section 2 by presenting a number of preliminary results, which we will need for the proofs of our main theorems. For example, the probabilistic technique highlighted above is introduced in Section 2.1, and we provide a brief discussion of some of our main computational methods in Section 2.4 (referring the reader to the supplementary file [1] for more details). Our proof of Theorem 1 for S_n and A_n is given in Section 3, where the details are divided into two subsections. In Section 3.1 we focus on the tuples (H_1, \dots, H_k) , where each H_i acts primitively on $\{1, \dots, n\}$, using fixed point ratio estimates to prove parts (ii) and (iii) of Theorem 1. These results feed in to our proof of Theorem 1(i) in Section 3.2, which requires an in-depth analysis of the tuples involving intransitive and imprimitive subgroups. Finally, we present our proof of Theorem 2 in Section 4, handling parts (i) and (ii) in Sections 4.1 and 4.2, respectively.

Acknowledgements. We thank Thomas Breuer, Derek Holt, Jürgen Müller and Eamonn O'Brien for their generous assistance with some of the computations involving sporadic groups. We also thank Peter Cameron for helpful comments and for drawing our attention to his paper [19]. The first author thanks the Heilbronn Institute for Mathematical Research for providing financial support during her doctoral studies at the University of Bristol.

2. PRELIMINARIES

In this section, we record several preliminary results that we will need in the proofs of our main theorems. More precisely, in Section 2.1 we adapt a probabilistic approach due to Liebeck and Shalev [36] for bounding the base size of a finite permutation group, which gives a useful technique for determining if a given subgroup tuple is regular. In Section 2.2, we highlight a connection between the base and independence numbers of a finite group, which was first observed by Cameron [19], and we use this to compute the base numbers $B(S_n)$ and $B(A_n)$. Next, in Section 2.3 we record several results that apply in the special case where G is almost simple. This includes Proposition 2.12, which shows that there are infinitely many finite simple groups with $B(G) < R(G)$. Finally, in Section 2.4 we briefly describe some of the main computational methods we employ in this paper, with further details provided in the supplementary file [1].

2.1. Probabilistic methods. Recall that if $G \leq \text{Sym}(\Omega)$ is a transitive permutation group with point stabiliser H , then

$$\text{fpr}(x, \Omega) = \frac{|C_\Omega(x)|}{|\Omega|} = \frac{|x^G \cap H|}{|x^G|}$$

is the *fixed point ratio* of $x \in G$, where $C_\Omega(x) = \{\alpha \in \Omega : \alpha^x = \alpha\}$ is the set of fixed points of x and x^G is the conjugacy class of x in G . Note that $\text{fpr}(x, \Omega)$ is the probability that x fixes a randomly chosen element of Ω (with respect to the uniform distribution on Ω).

In [36], Liebeck and Shalev introduced a powerful probabilistic method, based on fixed point ratio estimates, for bounding the base size of a finite permutation group. They used this approach to resolve the Cameron-Kantor base size conjecture from [21], which asserts that there is an absolute constant c such that $B_{\text{ns}}(G) \leq c$ for every almost simple group G (here $B_{\text{ns}}(G)$, as defined in Section 1, is the maximal base size $b(G, H)$ over all *non-standard* maximal subgroups H of G). The same approach was at the heart of the proof of Cameron's conjecture in the sequence of papers [7, 14, 15, 16], which established the optimal bound $B_{\text{ns}}(G) \leq 7$ in this setting.

The following key lemma is a natural generalisation and it encapsulates our probabilistic approach for studying the regularity of subgroup tuples.

Lemma 2.1. *Let G be a finite group and let $\tau = (H_1, \dots, H_k)$ be a core-free tuple of subgroups of G . Then τ is regular if*

$$\widehat{Q}(G, \tau) := \sum_{i=1}^t |x_i^G| \cdot \left(\prod_{j=1}^k \text{fpr}(x_i, G/H_j) \right) < 1,$$

where x_1, \dots, x_t is a set of representatives of the conjugacy classes in G of elements of prime order.

Proof. Set $\Gamma_i = G/H_i$ and $\Gamma = \Gamma_1 \times \dots \times \Gamma_k$. Let

$$Q(G, \tau) = \frac{|\{(\alpha_1, \dots, \alpha_k) \in \Gamma : \bigcap_{i=1}^k G_{\alpha_i} \neq 1\}|}{|\Gamma|}$$

be the probability that a randomly chosen tuple in Γ is not contained in a regular orbit. Then τ is regular if and only if $Q(G, \tau) < 1$, and thus it suffices to show that $Q(G, \tau) \leq \widehat{Q}(G, \tau)$.

A k -tuple $(\alpha_1, \dots, \alpha_k) \in \Gamma$ is not in a regular G -orbit if and only if there exists an element $x \in G$ of prime order such that $\alpha_i^x = \alpha_i$ for all i . Now the probability that an element $x \in G$ fixes a random element of Γ is equal to

$$\frac{|\{(\alpha_1, \dots, \alpha_k) \in \Gamma : \alpha_i^x = \alpha_i \text{ for all } i\}|}{|\Gamma|} = |\Gamma|^{-1} \prod_{j=1}^k |C_{\Gamma_j}(x)| = \prod_{j=1}^k \text{fpr}(x, \Gamma_j)$$

and thus

$$Q(G, \tau) \leq \sum_{x \in \mathcal{P}} \left(\prod_{j=1}^k \text{fpr}(x, \Gamma_j) \right) = \sum_{i=1}^t |x_i^G| \cdot \left(\prod_{j=1}^k \text{fpr}(x_i, \Gamma_j) \right) = \widehat{Q}(G, \tau),$$

where $\mathcal{P} = \bigsqcup_{i=1}^t x_i^G$ is the set of prime order elements of G . The result follows. \square

The following elementary result generalises [7, Lemma 2.1]. It provides a useful way to produce an upper bound on the expression $\widehat{Q}(G, \tau)$ in Lemma 2.1.

Lemma 2.2. *Let G be a finite group and let $\tau = (H_1, \dots, H_k)$ be a core-free tuple of subgroups of G . Suppose x_1, \dots, x_m are elements in G such that $|x_i^G| \geq B$ for all i and $\sum_{i=1}^m |x_i^G \cap H_j| \leq A_j$ for all j . Then*

$$\sum_{i=1}^m |x_i^G| \cdot \left(\prod_{j=1}^k \text{fpr}(x_i, G/H_j) \right) \leq B^{1-k} \cdot \prod_{j=1}^k A_j$$

Proof. We have

$$\sum_{i=1}^m |x_i^G| \cdot \left(\prod_{j=1}^k \text{fpr}(x_i, G/H_j) \right) = \sum_{i=1}^m |x_i^G|^{1-k} \cdot \left(\prod_{j=1}^k |x_i^G \cap H_j| \right) \leq B^{1-k} \cdot \left(\sum_{i=1}^m \prod_{j=1}^k |x_i^G \cap H_j| \right)$$

and thus

$$\sum_{i=1}^m |x_i^G| \cdot \left(\prod_{j=1}^k \text{fpr}(x_i, G/H_j) \right) \leq B^{1-k} \cdot \left(\prod_{j=1}^k \sum_{i=1}^m |x_i^G \cap H_j| \right) \leq B^{1-k} \cdot \prod_{j=1}^k A_j,$$

as claimed. \square

2.2. Independent sets. A subset S of a finite group G is *independent* if none of its elements lie in the subgroup generated by the others (that is to say, $x \notin \langle S \setminus \{x\} \rangle$ for all $x \in S$). For example, $\{(1, 2), (2, 3), \dots, (n-1, n)\}$ is an independent generating set for $G = S_n$.

We define the *independence number* of G , denoted $\sigma(G)$, to be the maximal size of an independent subset of G .

Proposition 2.3. *We have $B(G) \leq \sigma(G)$ for every finite group G .*

Proof. Suppose G acts faithfully on a set Ω and let $\mathcal{B} = \{\alpha_1, \dots, \alpha_b\}$ be a minimal base for G (in the sense that no proper subset of \mathcal{B} is a base). In [19], Cameron reveals a beautiful connection with lattices in order to establish the bound $b \leq \sigma(G)$ (see [19, Proposition 3.2]). We can also argue directly as follows.

For all $i \in \{1, \dots, b\} = [b]$, let

$$H_i = \bigcap_{j \in [b] \setminus \{i\}} G_{\alpha_j}$$

and note that the minimality of \mathcal{B} implies that each H_i is nontrivial. Fix a nontrivial element $g_i \in H_i$ and observe that $g_i \notin G_{\alpha_i}$. Then the g_i are distinct and we claim that $\{g_1, \dots, g_b\}$ is an independent subset of G . To see this, set $K_i = \langle g_1, \dots, g_{i-1}, g_{i+1}, \dots, g_b \rangle$ for all $i \in [b]$. Since $g_j \in G_{\alpha_i}$ for all $j \neq i$, we have $K_i \leq G_{\alpha_i}$ for all i . But $g_i \notin G_{\alpha_i}$ and thus $g_i \notin K_i$. This justifies the claim and we conclude that $b \leq \sigma(G)$.

In particular, $B(G) \leq \sigma(G)$, as required. \square

We can now combine this observation with a theorem of Whiston [53] to compute the base numbers of S_n and A_n (also see [19, Corollary 3.4]).

Corollary 2.4. *For $n \geq 5$ we have $B(S_n) = n - 1$ and $B(A_n) = n - 2$.*

Proof. By considering the natural action of S_n and A_n of degree n , we see that $B(S_n) \geq n - 1$ and $B(A_n) \geq n - 2$. We now combine this trivial observation with the main theorem of [53], which states that every independent subset in S_n has at most $n - 1$ elements, with equality only if the subset generates S_n (for example, $\{(1, 2), (2, 3), \dots, (n-1, n)\}$ is an independent subset of maximal size). It follows that $\sigma(S_n) = n - 1$ and $\sigma(A_n) \leq n - 2$, and we conclude by applying Proposition 2.3. \square

Remark 2.5. Let $G = S_n$ or A_n , and let H be a core-free maximal subgroup of G . Through the combined efforts of several authors, the base size $b(G, H)$ is now known in all cases:

- (a) H primitive on $\{1, \dots, n\}$: See [14]
- (b) H imprimitive: See [12, 33, 43]
- (c) H intransitive: See [24, 31, 42]

2.3. Almost simple groups. Let G be a finite almost simple group with socle T , which means that

$$T \trianglelefteq G \leq \text{Aut}(T),$$

where we identify T with its group of inner automorphisms. Let us define the regularity and base number of G , denoted $R(G)$ and $B(G)$ respectively, as in Section 1, and recall that $R(G) \geq B(G)$. We begin by recording the following trivial observation, which applies in the general setting.

Lemma 2.6. *Let G be a finite group and let $\tau = (H_1, \dots, H_k)$ be a k -tuple of core-free subgroups of G . Then τ is non-regular if*

$$\prod_{i=1}^k |H_i| > |G|^{k-1}.$$

Proof. Clearly, G has a regular orbit on $G/H_1 \times \dots \times G/H_k$ only if

$$|G| \leq \prod_{i=1}^k |G : H_i|$$

and the result follows. \square

Proposition 2.7. *If G is a finite almost simple group, then $B(G) \geq 3$.*

Proof. By [13, Proposition 2.5], G has a core-free subgroup H with $|H|^2 > |G|$ and thus (H, H) is a non-regular pair by Lemma 2.6. \square

It will be useful to introduce the following definition.

Definition 2.8. Let G be an almost simple group with socle T . We define $\mathcal{M}(G)$ to be the set of core-free subgroups of G that are maximal in some overgroup of T .

For example, $H = \text{AGL}_3(2) \in \mathcal{M}(S_8)$ since it is a maximal subgroup of A_8 (and note that H is not contained in a core-free maximal subgroup of S_8). This definition allows us to present the following elementary lemma, which tells us that in the almost simple setting, we only need to consider tuples of subgroups in $\mathcal{M}(G)$ in order to bound $R(G)$ from above.

Lemma 2.9. *Let G be an almost simple group with socle T . Then $R(G) \leq k$ if and only if every k -tuple of subgroups in $\mathcal{M}(G)$ is regular.*

Proof. The forward implication is clear, so let us assume every k -tuple of subgroups in $\mathcal{M}(G)$ is regular and let $\tau = (H_1, \dots, H_k)$ be an arbitrary core-free tuple. Then each H_i is contained in a subgroup $M_i \in \mathcal{M}(G)$ and the regularity of (M_1, \dots, M_k) immediately implies that τ is also regular. \square

Remark 2.10. Notice that $|G : T| \leq 2$ for every almost simple group G we are considering in Theorems 1 and 2, with the single exception of the special case $G = A_6.2^2$. If $G = T.2$ and $H < T$ is maximal, then H is contained in a core-free maximal subgroup of G if $N_G(H) \neq H$. So it follows that $R(G) \leq k$ if and only if every k -tuple of subgroups in $\mathcal{M}'(G)$ is regular, where the latter comprises the core-free maximal subgroups of G , together with any maximal subgroups H of T with $N_G(H) = H$. This minor refinement of Lemma 2.9 will be useful in the proof of Theorem 2 (see the discussion at the start of Section 4.1).

The following easy observation will be useful when we study soluble tuples for sporadic groups in Section 4.2.

Lemma 2.11. *Let G be a finite group and let H be a simple subgroup of G with $R_{\text{sol}}(H) \leq k$. Then every core-free soluble k -tuple of subgroups of G of the form $\tau = (H_1, \dots, H_k)$ with $H_1 < H$ is regular.*

Proof. Since $R_{\text{sol}}(H) \leq k$, there exist $g_i \in H$ such that

$$1 = H_1^{g_1} \cap \bigcap_{i=2}^k (H_i \cap H)^{g_i} = \bigcap_{i=1}^k H_i^{g_i}$$

and the result follows. \square

As noted in Remark 1(a), there are examples where the trivial bound $B(G) \leq R(G)$ is a strict inequality. In fact, we can show that there are infinitely many simple groups with this property.

Proposition 2.12. *There are infinitely many finite simple groups G with $B(G) < R(G)$.*

Proof. Let $G = \text{SL}_3(q)$, where $q \geq 3$ and $q \not\equiv 1 \pmod{3}$. Let V be the natural module and let $H = P_1$ be the stabiliser in G of a 1-dimensional subspace of V . Similarly, let K be the stabiliser of a 2-space. Then $b(G, H) = b(G, K) = 4$ and by applying the main theorem of [7] we deduce that $B(G) = 4$. So in order to prove the proposition, it suffices to establish the following claim:

Claim. *(H, H, K, K) is a non-regular 4-tuple.*

To see this, let $\langle v_1 \rangle$ and $\langle v_2 \rangle$ be distinct 1-spaces and choose v_3 so that $\beta = \{v_1, v_2, v_3\}$ is a basis for V . Let H_i be the stabiliser of $\langle v_i \rangle$ and let $L = H_1 \cap H_2$, so $|L| = q^2(q-1)^2$. Set $\Omega = G/K$ and identify Ω with the set of 2-dimensional subspaces of V . It suffices to show that $b(L, \Omega) \geq 3$. Equivalently, we need to show that no point stabiliser L_α (with $\alpha \in \Omega$) has a regular orbit on Ω .

Set $U_1 = \langle v_1, v_2 \rangle$, $U_2 = \langle v_1, v_3 \rangle$ and $U_3 = \langle v_2, v_3 \rangle$. Then these 2-spaces represent three distinct L -orbits, with respective lengths 1, q and q . In addition, the stabiliser in L of $U_4 = \langle v_1 + v_2, v_3 \rangle$ is the subgroup

$$L_4 = \left\{ \begin{pmatrix} a & 0 & b \\ 0 & a & b \\ 0 & 0 & a^{-2} \end{pmatrix} : a \in \mathbb{F}_q^\times, b \in \mathbb{F}_q \right\}$$

with respect to the basis β , whence the L -orbit of U_4 has size $q(q-1)$. Since $|\Omega| = q^2 + q + 1$, we deduce that L has exactly 4 orbits on Ω , with respective stabilisers $L_i = L_{U_i}$ for $i = 1, 2, 3, 4$.

So to prove the claim, it just remains to show that none of these point stabilisers has a regular orbit on Ω . This is clear for $i = 1, 2, 3$ since $|L_i| > |\Omega|$. So let us consider L_4 . Clearly, the L_4 -orbits of U_1 and U_4 both have length 1, and it is straightforward to show that U_2 and U_3 lie in distinct L_4 -orbits of length q . So these four orbits have already covered $2q + 2$ points in Ω and we note that $|\Omega| - (2q + 2) < |L_4|$. Therefore, L_4 does not have a regular orbit on Ω and the proof of the claim (and proposition) is complete. \square

The previous proposition shows that there are infinitely many finite simple groups with $B(G) < R(G)$. This observation is extended in our next result, which demonstrates the existence of simple groups G for which $R(G) - B(G)$ can be arbitrarily large.

Proposition 2.13. *Let $G = \text{L}_n(2)$ with $n \geq 5$. Then $B(G) = n$ and $R(G) \geq 2(n-1)$.*

Proof. First observe that $G = \text{GL}(V)$, where V is an n -dimensional vector space over \mathbb{F}_2 .

Let H be a maximal subgroup of G . If H acts irreducibly on V , then the main theorem of [7] implies that $b(G, H) \leq 4$. Now assume H is reducible, which means that we can identify G/H with the set Ω of k -dimensional subspaces of V for some positive integer $k \leq n/2$. Clearly, if $k = 1$ then Ω coincides with the set of nonzero vectors in V , in which case a subset of Ω is a base if and only if it is a basis and thus $b(G, H) = n$. So in order to prove that $B(G) = n$, it suffices to show that $b(G, H) \leq n$ for all $2 \leq k \leq n/2$. To see this, fix a basis

$\{e_1, \dots, e_n\}$ for V and define the following k -dimensional subspaces of V :

$$\begin{aligned} V_1 &= \langle e_1, e_2, \dots, e_k \rangle \\ V_2 &= \langle e_2, e_3, \dots, e_{k+1} \rangle \\ &\vdots \\ V_n &= \langle e_n, e_1, \dots, e_{k-1} \rangle \end{aligned}$$

Let L_i be the setwise stabiliser of V_i in G and set $L = \bigcap_i L_i$. Now each basis vector e_i is contained in exactly k of these subspaces, say V_{a_1}, \dots, V_{a_k} , and one checks that $\bigcap_j V_{a_j} = \langle e_i \rangle$. Since we are working over \mathbb{F}_2 , it follows that L fixes each e_i and thus $L = 1$. In other words, $\{V_1, \dots, V_n\}$ is a base for G and we conclude that $b(G, H) \leq n$, as required.

Let H and K be the stabilisers of a 1-space and an $(n-1)$ -space, respectively. In order to show that $R(G) \geq 2(n-1)$, it suffices to prove the following claim.

Claim. *The $(2n-3)$ -tuple $\tau = (H, \dots, H, K, \dots, K)$ comprising $n-1$ copies of H and $n-2$ copies of K is non-regular.*

To see this, let g_1, \dots, g_{n-1} be arbitrary elements in G and set $L = \bigcap_i H^{g_i}$. Here H^{g_i} is the stabiliser of a nonzero vector $v_i \in V$ and L contains the following elementary abelian 2-subgroup (with respect to an appropriate basis for V)

$$J = \left\{ \left(\begin{array}{c|c} I_{n-1} & A \\ \hline 0 & I_1 \end{array} \right) : A \in \mathbb{F}_2^{n-1} \right\}$$

Note that $L = J$ if the v_i are linearly independent and we work with a basis of the form $\{v_1, \dots, v_{n-1}, v_n\}$ for V .

So to prove the claim, it suffices to show that $b(J, \Omega) \geq n-1$, where J is defined as above (with respect to a basis $\{v_1, \dots, v_{n-1}, v_n\}$ for V) and Ω is the set of $(n-1)$ -dimensional subspaces of V . Notice that if τ denotes the inverse-transpose graph automorphism of G , then we have $b(J, \Omega) = b(J^\tau, \Gamma)$, where Γ is the set of nonzero vectors in V , so we need to show that $b(J^\tau, \Gamma) \geq n-1$.

Let w_1, \dots, w_{n-2} be arbitrary nonzero vectors in V and write

$$w_i = b_{i,1}v_1 + \dots + b_{i,n-1}v_{n-1} + b_{i,n}v_n$$

for all i . Let $B = (b_{i,j})$ be the $(n-2) \times (n-1)$ matrix corresponding to the coefficients of v_j for $j < n$. Then the pointwise stabiliser of the w_i in J^τ is the set of matrices of the form

$$\left(\begin{array}{c|c} I_{n-1} & 0 \\ \hline A & I_1 \end{array} \right) \in J^\tau$$

where $A = (a_1, \dots, a_{n-1}) \in \mathbb{F}_2^{n-1}$ and $BA^T = 0$. Since the kernel of the linear map $\mathbb{F}_2^{n-1} \rightarrow \mathbb{F}_2^{n-2}$ corresponding to B is obviously nonzero, it follows that the pointwise stabiliser of the w_i in J^τ is nontrivial. In turn, this implies that $b(J^\tau, \Gamma) = b(J, \Omega) \geq n-1$ and thus τ is non-regular as claimed. \square

Remark 2.14.

- (a) Notice that in Proposition 2.13 we assume $G = L_n(2)$ with $n \geq 5$. For $n = 3$ we get $B(G) = 3$ and $R(G) = 4$. However, the case $n = 4$ is different since $L_4(2) \cong A_8$ and thus $B(G) = R(G) = 6$ (as a special case of Theorem 1(i)).
- (b) We expect $R(G) = 2(n-1)$ when $G = L_n(2)$ and $n \geq 3$, which would imply that $R(G) < 2B(G)$. In particular, this would be consistent with Conjecture 1(i), which asserts (as a special case) that there is an absolute constant c such that $R(G) < c \cdot B(G)$ for every finite simple group G .

2.4. Computational methods. To conclude this preliminary section, we briefly discuss some of the main computational methods we apply in this paper. All of our computations are performed using MAGMA [5] (version 2.28-4) and GAP [28] (version 4.11.1). We refer the reader to the supplementary file [1] for further details, including sample code.

Let G be a finite group and let $\tau = (H_1, \dots, H_k)$ be a k -tuple of core-free subgroups with $k \geq 2$. As noted in Lemma 2.6, τ is non-regular if $\prod_i |H_i| > |G|^{k-1}$.

2.4.1. Random search. If a suitable permutation or matrix representation of G is available, then random search often provides an efficient way to show that τ is regular. Here the aim is to find elements $g_i \in G$, in terms of the given representation, such that

$$\bigcap_{i=1}^k H_i^{g_i} = 1.$$

Typically, we are interested in the case where τ is a maximal tuple, which means that each component H_i is a core-free maximal subgroup of G .

Of course, the groups S_n and A_n are naturally defined in terms of a permutation representation. For an almost simple sporadic group $G \notin \{J_4, \text{Ly}, \text{Th}, \mathbb{B}, \mathbb{M}\}$, we use the MAGMA function `AutomorphismGroupSimpleGroup` to construct G as a permutation group on n points, where $n = m(G)$ is the minimal degree of a faithful permutation representation. Similarly, for $G \in \{J_4, \text{Ly}, \text{Th}\}$ we can use the function `MatrixGroup` to construct G as a matrix group. For example, this allows us to view the Lyons group as a subgroup of $\text{GL}_{111}(5)$. However, the largest sporadic groups \mathbb{B} and \mathbb{M} do not admit a permutation or matrix representation that is suitable for direct computation, so they will require a different approach (for example, $97239461142009186000 \sim 9.7 \times 10^{19}$ is the minimal degree of a faithful permutation representation of \mathbb{M}).

Given a suitable representation of G , we can use the MAGMA function `MaximalSubgroups` to construct a complete set of representatives of the conjugacy classes of maximal subgroups of G . For example, this is effective for all $G \in \{S_n, A_n\}$ with $n \leq 200$. And similarly for all sporadic groups G , unless $G \in \{\text{Fi}'_{24}, \text{Fi}_{24}, \text{Th}\}$. For the latter groups, we can construct the maximal subgroups we will need to work with via explicit generators presented in the Web Atlas [49], which are given as words in the standard generators for G (these generators can also be constructed using the GAP package `AtlasRep` [52]). We can of course filter the output from `MaximalSubgroups` if we are just interested in soluble maximal subgroups, or the maximal subgroups of S_n that act primitively on $\{1, \dots, n\}$, etc.

For some of the groups we are interested in, we can replace `MaximalSubgroups` by other MAGMA functions, such as `SolubleSubgroups` and `NilpotentSubgroups`, which return a complete set of representatives of the conjugacy classes of soluble (respectively, nilpotent) subgroups of G (not surprisingly, the effectiveness of the latter functions is more limited for groups of large order). We can also iteratively apply `MaximalSubgroups` in order to descend deeper into the subgroup lattice of G . For example, this will be a useful technique in the proof of Theorem 2(ii), when we are seeking to show that $R_{\text{sol}}(G) \leq 3$ in the setting where G is a sporadic group and the function `SolubleSubgroups` is unavailable (see the proofs of Lemmas 4.16 and 4.17, for instance).

2.4.2. Orbit computations. Suppose we can construct G and each H_i , in terms of a suitable permutation or matrix representation, and let us assume random search is inconclusive. Here it is useful to observe that τ is regular if and only if H_1 has a regular orbit on

$$Y = G/H_2 \times \cdots \times G/H_k.$$

So assuming that the indices $|G : H_i|$ are not prohibitively large, we can use the MAGMA function `CoSetAction` to construct the action on G on each set G/H_i with $i \geq 2$, which will allow us to calculate all of the orbits of H_1 on Y and then determine whether or not τ is

regular. Note that in order to conclude that τ is non-regular, it suffices to find a collection of distinct non-regular H_1 -orbits Y_1, \dots, Y_t such that $\sum_i |Y_i| > |Y| - |H_1|$.

In the special case $k = 2$, we can also use `DoubleCosetRepresentatives` to calculate the size of every (H_1, H_2) double coset in G , noting that τ is regular if and only if there exists a double coset of size $|H_1||H_2|$.

Remark 2.15. There are a handful of maximal pairs $\tau = (H_1, H_2)$ for $G \in \{J_4, L_3, Th\}$ where we have $|H_1||H_2| < |G|$, random search is inconclusive and both of the above methods for determining whether or not τ is regular are ineffective due to the very large size of the indices $|G : H_1|$ and $|G : H_2|$. We thank Derek Holt, Jürgen Müller and Eamonn O'Brien for resolving these special cases, which are presented in Lemma 4.8. In particular, Müller used the GAP package ORB [44] to show that the maximal pair for $G = J_4$ with

$$H_1 = 2^{3+12}.(S_5 \times L_3(2)), \quad H_2 = 2^{1+12}.3.M_{22}:2$$

is non-regular (here we have $|G : H_1| = 131358148251$ and $|G : H_2| = 3980549947$).

2.4.3. Character theory. If we have access to the character tables of G and each component subgroup H_i , together with the fusion maps from H_i -classes to G -classes, then we can compute

$$\text{fpr}(x, G/H_i) = \frac{|x^G \cap H_i|}{|x^G|}$$

precisely for all $x \in G$ and all i . In turn, this allows us to evaluate the expression $\widehat{Q}(G, \tau)$ in Lemma 2.1, recalling that τ is regular if $\widehat{Q}(G, \tau) < 1$. This can be an effective way to produce a list of candidate non-regular k -tuples, which can then be analysed using other methods (such as random search, as described above).

The character table of every almost simple sporadic group G is available in the GAP Character Table Library [6]. In addition, if $G \neq \mathbb{M}$, then the character table of every maximal subgroup of G is also available in [6] and we can use the GAP function `Maxes` to access the associated fusion maps on conjugacy classes. For $G \neq \mathbb{M}$, this allows us to compute $\widehat{Q}(G, \tau)$ for every maximal tuple τ (if $G = \mathbb{M}$, we can handle some, but not all, maximal tuples in the same way, using the function `NamesOfFusionSources` to access the relevant character tables and fusion maps). We will rely heavily on this probabilistic approach to prove Theorem 2 for the large sporadic groups, including \mathbb{B} and \mathbb{M} (see the proofs of Lemmas 4.7, 4.9 and 4.10, for example).

3. SYMMETRIC AND ALTERNATING GROUPS

In this section we assume G is an almost simple group with socle A_n and our aim is to prove Theorem 1. We begin in Section 3.1 by considering parts (ii) and (iii), focussing initially on primitive tuples (indeed, part (iii) quickly follows from part (ii) since every soluble maximal subgroup of G is primitive when $n \geq 17$). And then in Section 3.2 we turn our attention to the regularity number, proving part (i) of Theorem 1. To do this, we first handle the tuples where each subgroup acts intransitively on $\{1, \dots, n\}$ (see Section 3.2.1). The imprimitive tuples are then treated in Section 3.2.2 and we bring everything together in Section 3.2.3, where we complete the proof of Theorem 1.

3.1. Primitive subgroups. Let G be an almost simple group with socle $T = A_n$ and recall that a subgroup H of G is said to be *primitive* if $H \cap T$ acts primitively on the set $[n] = \{1, \dots, n\}$. We define $R_{\text{prim}}(G)$ to be the minimal positive integer k such that every core-free primitive k -tuple of subgroups of G is regular. Similarly, $B_{\text{prim}}(G)$ is the maximal base size $b(G, H)$ over all core-free primitive subgroups H of G . Of course, if G does not have a core-free primitive subgroup, then $R_{\text{prim}}(G)$ and $B_{\text{prim}}(G)$ are undefined.

In this section we will prove parts (ii) and (iii) of Theorem 1. As a consequence, we establish a strong form of Conjecture 1(ii) for symmetric and alternating groups, and we take the first steps towards Conjecture 1(iii) in this setting.

$R_{\text{prim}}(G)$	G	τ
6	A_8	$(\text{AGL}_3(2)_1, \text{AGL}_3(2)_1, \text{AGL}_3(2)_1, \text{AGL}_3(2)_2, \text{AGL}_3(2)_2)$
5	S_6	$(\text{PGL}_2(5), \text{PGL}_2(5), \text{PGL}_2(5), \text{PGL}_2(5))$
4	A_6	$(\text{L}_2(5), \text{L}_2(5), \text{L}_2(5))$
	$A_6.2 \neq S_6$	$(\text{L}_2(5), \text{L}_2(5), \text{L}_2(5))$
	$A_6.2^2$	$(\text{L}_2(5), \text{L}_2(5), \text{L}_2(5))$
	A_7	$(\text{L}_2(7)_1, \text{L}_2(7)_1, \text{L}_2(7)_2)$
	S_8	$(\text{PGL}_2(7), \text{PGL}_2(7), \text{PGL}_2(7))$
3	A_5	$(\text{D}_{10}, \text{D}_{10})$
	S_5	$(\text{AGL}_1(5), \text{AGL}_1(5))$
	S_7	$(\text{L}_2(7), \text{L}_2(7))$
	A_9	$(\text{P}\Gamma\text{L}_2(8), \text{P}\Gamma\text{L}_2(8))$
	S_9	$(\text{AGL}_2(3), \text{AGL}_2(3))$
	S_{10}	$(\text{P}\Gamma\text{L}_2(9), \text{P}\Gamma\text{L}_2(9))$
	A_{11}	$(\text{M}_{11}, \text{M}_{11})$
	S_{11}	$(\text{M}_{11}, \text{M}_{11})$
	A_{12}	$(\text{M}_{12}, \text{M}_{12})$
	S_{12}	$(\text{M}_{12}, \text{M}_{12})$

TABLE 1. Almost simple groups G with socle A_n and $R_{\text{prim}}(G) > 2$

Let us briefly outline our approach. We begin by handling the groups of small degree in Proposition 3.1 below, which are amenable to direct computation in MAGMA [5]. Our method for dealing with the general case relies on the probabilistic approach described in Section 2.1. To do this, we will apply two key results in order to derive the required fixed point ratio estimates, which we combine in the statement of Theorem 3.3. The first is a theorem of Maróti [40] on the orders of core-free primitive subgroups of S_n . The second main ingredient is a theorem of Guralnick and Magaard [30] on the minimal degree of primitive subgroups, which we can use to obtain lower bounds on the conjugacy class sizes of prime order elements lying in a primitive subgroup. Our approach is similar to the one adopted in [14], where the main theorem states that $B_{\text{prim}}(G) \leq 5$, with $B_{\text{prim}}(G) = 2$ if $n \geq 13$.

In order to establish the bound $R_{\text{prim}}(G) \leq k$, it suffices to show that every k -tuple of primitive subgroups in $\mathcal{M}(G)$ is regular, where $\mathcal{M}(G)$ is the set of core-free subgroups that are maximal in some subgroup of G containing T (see Definition 2.8 and Lemma 2.9). In fact, if $G = S_n$ or A_n then we only need to consider tuples involving subgroups in $\mathcal{M}'(G)$, which is the set of core-free maximal subgroups of G , together with any maximal subgroups H of T with $N_G(H) = H$ (see Remark 2.10).

We begin by establishing part (ii) of Theorem 1 for the groups with $n < 60$.

Proposition 3.1. *Let G be an almost simple group with socle $T = A_n$, where $n < 60$, and assume G has a core-free primitive subgroup.*

- (i) *We have $R_{\text{prim}}(G) \leq 6$, with equality if and only if $G = A_8$.*
- (ii) *If $n \geq 13$, then $R_{\text{prim}}(G) = 2$. For $n \leq 12$, the groups with $R_{\text{prim}}(G) = k > 2$ are recorded in Table 1, together with a non-regular primitive tuple $\tau = (H_1, \dots, H_{k-1})$.*

Proof. This is a routine MAGMA [5] computation. For $n \geq 13$, we can use random search (see Section 2.4.1) to show that every primitive pair (H_1, H_2) with $H_i \in \mathcal{M}'(G)$ is regular. Similarly, for $n \leq 12$ it is straightforward to identify all the non-regular pairs (H_1, H_2) with $H_i \in \mathcal{M}(G)$, which we can then use to determine any non-regular triples with components in $\mathcal{M}(G)$. If required, we can then extend the analysis to 4-tuples and 5-tuples, which allows us to deduce that every primitive 6-tuple is regular. \square

Remark 3.2. We record some remarks on the statement of Proposition 3.1.

- (a) Notice that if $n \leq 12$, then $G = A_{10}$ is the only group with $R_{\text{prim}}(G) = 2$. Here $H = \text{M}_{10}$ represents the unique class of primitive maximal subgroups of G and one checks that $b(G, H) = 2$, so every primitive pair for G is regular.

- (b) By inspecting [14], we deduce that if $n < 60$ then $R_{\text{prim}}(G) = B_{\text{prim}}(G)$ unless $G = A_7$ or A_8 , where we have

$$\begin{aligned} R_{\text{prim}}(A_7) &= B_{\text{prim}}(A_7) + 1 = 4 \\ R_{\text{prim}}(A_8) &= B_{\text{prim}}(A_8) + 2 = 6 \end{aligned}$$

- (c) The group $G = A_8$ has two conjugacy classes of maximal subgroups isomorphic to $\text{AGL}_3(2)$, represented by H and K . Here $b(G, H) = b(G, K) = 4$, but we find that the 5-tuple (H, H, H, K, K) is non-regular, as indicated in the first row of Table 1 (we use subscripts 1 and 2 to distinguish representatives of the two classes).
- (d) Similarly, $G = A_7$ has two classes of maximal subgroups isomorphic to $L_2(7)$ and (H, H, K) is a non-regular triple, where H and K represent these two conjugacy classes (note that $b(G, H) = b(G, K) = 3$).

In order to complete the proof of Theorem 1(ii), we may assume $n \geq 60$. So $G = S_n$ or A_n , and our goal is to prove that $R_{\text{prim}}(G) = 2$.

Recall that the *minimal degree* of a subgroup H of G , denoted $\mu(H)$, is the minimal number of points in $[n] = \{1, \dots, n\}$ moved by a non-identity element of H . That is,

$$\mu(H) = \min\{\text{supp}(x) : 1 \neq x \in H\}$$

where $\text{supp}(x)$, the *support* of x , is the number of points moved by x .

As explained above, we will complete the proof of Theorem 1(iii) by applying the probabilistic approach encapsulated in Lemma 2.1, which relies on key theorems of Maróti [40] and Guralnick and Magaard [30]. We combine (and slightly simplify) both results in the following statement.

Theorem 3.3. *Suppose $n \geq 60$ and $H \in \mathcal{M}(G)$ is primitive. Then either*

$$|H| < n^{1+\lceil \log_2 n \rceil} \text{ and } \mu(H) \geq n/2 - \sqrt{n},$$

or one of the following holds:

- (i) $H = (S_l \wr S_k) \cap G$ in its product action on $n = l^k$ points, where $l \geq 5$ and $k \geq 2$, and we have $\mu(H) = 2l^{k-1}$.
- (ii) $H = S_l \cap G$ in its action on the k -element subsets of $\{1, \dots, l\}$, so $n = \binom{l}{k}$ with $l \geq 5$ and $2 \leq k < l/2$, and we have $\mu(H) = 2 \binom{l-2}{k-1}$.

Proof. The bound on $|H|$ follows immediately from [40, Theorem 1.1]. Now let us turn to the minimal degree of H . By [30, Theorem 1], if we exclude the cases recorded in (i) and (ii), then either $\mu(H) \geq n/2$, or $H = \text{O}(V)$ is an orthogonal group over the field \mathbb{F}_2 acting on a set of hyperplanes of the natural module V . In each of the latter cases, the exact value of $\mu(H)$ is given in the statement of [30, Theorem 1] and it is easy to check that the desired bound $\mu(H) \geq n/2 - \sqrt{n}$ is satisfied. \square

Next we seek lower bounds on the size of a conjugacy class x^G defined in terms of the support of x . This will allow us to translate the minimal degree bound in Theorem 3.3 into a lower bound on $|x^G|$, which applies whenever x is a prime order element contained in an appropriate primitive subgroup of G . The following result holds for all $n \geq 5$.

Lemma 3.4. *Let $G = S_n$ or A_n with $n \geq 5$ and suppose $x \in G$ has prime order r with $\text{supp}(x) = m \geq \sqrt{n}$. Then $|x^G| \geq f_2(m)$, with $|x^G| \geq f_3(m)$ if r is odd, where*

$$f_s(m) = \frac{n!}{s^{m/s} [m/s]! (n-m)!}.$$

Proof. First observe that we may assume $G = S_n$. To see this, suppose $G = A_n$ and recall that $|x^G| = |x^{S_n}|$ unless x is an r -cycle and $r \in \{n-1, n\}$. If $r = n$ then $m = n$ and the

desired conclusion holds since

$$|x^G| = \frac{1}{2}(n-1)! > \frac{n!}{3^{n/3}\lceil n/3 \rceil!} > \frac{n!}{2^{n/2}\lceil n/2 \rceil!}$$

And similarly if $r = n - 1$. So for the remainder, we will assume $G = S_n$.

Suppose x has cycle-type $(r^a, 1^{n-ar})$ for some $a \geq 1$. Then $\text{supp}(x) = ar$ and

$$|x^G| = \frac{n!}{r^a a! (n-ar)!}.$$

In particular, the result follows immediately if $r = 2$, so we may assume r is odd.

Suppose there exists an element $y \in G$ of prime order $q < r$ with $\text{supp}(y) = m$. Then it is easy to check that $q^{m/q}(m/q)! > r^{m/r}(m/r)!$ and thus $|x^G| > |y^G|$. In particular, if m is even then we immediately deduce that $|x^G| \geq f_2(m)$. And the same conclusion holds for m odd since

$$2^{m/2}((m+1)/2)! \geq r^{m/r}(m/r)!$$

for every odd prime r dividing m . Finally, in order to establish the bound $|x^G| \geq f_3(m)$ it suffices to show that

$$3^{m/3}\lceil m/3 \rceil! \geq r^{m/r}(m/r)!$$

and this is straightforward to verify. \square

We now bring the minimal degree bound in Theorem 3.3 in to play, which allows us to derive a lower bound on the conjugacy class sizes of elements of prime order contained in a primitive maximal subgroup of S_n or A_n . In the statement, we refer to the function $f_s(m)$ defined in Lemma 3.4.

Lemma 3.5. *Let $G = S_n$ or A_n with $n \geq 60$ and suppose $x \in H$ has prime order r , where $H \in \mathcal{M}(G)$ is primitive.*

- (i) *We have $|x^G| \geq f_2(\ell)$, with $|x^G| \geq f_3(\ell)$ if r is odd, where $\ell = \lceil 2\sqrt{n} \rceil$.*
- (ii) *If $\mu(H) \geq n/2 - \sqrt{n}$, then $|x^G| \geq f_2(\ell')$ with $\ell' = \lceil n/2 - \sqrt{n} \rceil$.*

Proof. Set $m = \text{supp}(x)$. By applying Theorem 3.3, we deduce that $\mu(H) \geq 2\sqrt{n}$, which in turn implies that $\lceil 2\sqrt{n} \rceil \leq m \leq n$. By Lemma 3.4, it follows that $|x^G| \geq f_2(m)$, with $|x^G| \geq f_3(m)$ if r is odd, and it is straightforward to show that both lower bounds are minimal when $m = \lceil 2\sqrt{n} \rceil$, which establishes (i). Part (ii) is entirely similar. \square

We will also need an upper bound on the number of involutions in a primitive subgroup $H \in \mathcal{M}(G)$, which is denoted by $i_2(H)$.

Lemma 3.6. *If $n \geq 5$, then $i_2(H) \leq 5^{(n-1)/6} \sqrt{|H|}$ for every subgroup H of S_n .*

Proof. First recall that $i_2(H) \leq \sqrt{k|H|}$, where k is the number of conjugacy classes in H (this standard bound is easily obtained by considering the Frobenius-Schur indicators of the complex irreducible characters of H). By the main theorem of [29] we have $k \leq 5^{(n-1)/3}$ and the result follows. \square

Corollary 3.7. *Let $G = S_n$ or A_n with $n \geq 60$ and suppose $H \in \mathcal{M}(G)$ is primitive as in part (i) or (ii) of Theorem 3.3. Then*

$$i_2(H) \leq \lceil \sqrt{n} \rceil! (1 + 5^{(\sqrt{n}-1)/6})^2.$$

Proof. First assume that $H = (S_l \wr S_k) \cap G$, where $n = l^k$, $l \geq 5$ and $k \geq 2$, and note that it suffices to bound $i_2(S_l \wr S_k)$. For $k = 2$, Lemma 3.6 implies that there are fewer than $(1 + 5^{(l-1)/6}(l!)^{1/2})^2$ involutions in the base group $S_l \times S_l$ of the wreath product, and there are exactly $l!$ additional involutions in H . Therefore,

$$i_2(H) \leq (1 + 5^{(l-1)/6}(l!)^{1/2})^2 + l! < l!(1 + 5^{(l-1)/6})^2.$$

And for $k \geq 3$ one can check that

$$i_2(H) \leq |S_l \wr S_k| = (k\sqrt{n})! k! \leq \lceil \sqrt{n} \rceil! (1 + 5^{(\sqrt{n}-1)/6})^2 = f(n).$$

Now assume $H = S_l \wr G$ and $n = \binom{l}{k}$, where $l \geq 5$ and $2 \leq k < l/2$. For $k = 2$, Lemma 3.6 implies that

$$i_2(H) \leq i_2(S_l) \leq 5^{(l-1)/6} (l!)^{1/2} < 5^{\sqrt{2n}/6} \lceil \sqrt{2n} + 1 \rceil!^{1/2} = g(n)$$

and it is easy to check $i_2(H) \leq |H| \leq l! < g(n)$ for $k \geq 3$ and $n \geq 60$. The result now follows since $g(n) \leq f(n)$ for all $n \geq 60$. \square

We are now in a position to prove Theorem 1(ii).

Proof of Theorem 1(ii). In view of Proposition 3.1, we may assume $n \geq 60$, so $G = S_n$ or A_n , and our goal is to prove that $R_{\text{prim}}(G) = 2$. To do this, it suffices to show that every primitive pair $\tau = (H_1, H_2)$ is regular, where $H_1, H_2 \in \mathcal{M}(G)$. Define

$$\widehat{Q}(G, \tau) = \sum_{i=1}^t \frac{|x_i^G \cap H_1| |x_i^G \cap H_2|}{|x_i^G|}$$

as in Lemma 2.1, where x_1, \dots, x_t is a set of representatives of the conjugacy classes in G of elements of prime order, and recall that τ is regular if $\widehat{Q}(G, \tau) < 1$.

First assume that $\mu(H_1) \geq n/2 - \sqrt{n}$. Then $|x_i^G \cap H_1| = 0$ if $\text{supp}(x_i) < n/2 - \sqrt{n}$ and thus

$$\widehat{Q}(G, \tau) = \sum_{i=1}^s \frac{|y_i^G \cap H_1| |y_i^G \cap H_2|}{|y_i^G|}$$

where y_1, \dots, y_s represent the classes of prime order elements with support at least $n/2 - \sqrt{n}$. By combining Lemmas 2.2 and 3.5(ii) with Maróti's bound in Theorem 3.3 on the orders of H_1 and H_2 , we deduce that

$$\widehat{Q}(G, \tau) \leq n^{2+2\lceil \log_2 n \rceil} \cdot \frac{2^{\ell/2} \lceil \ell/2 \rceil! (n - \ell)!}{n!},$$

where $\ell = \lceil n/2 - \sqrt{n} \rceil$. One checks that this upper bound is less than 1 for all $n \geq 60$.

By the same argument, τ is regular if $\mu(H_2) \geq n/2 - \sqrt{n}$, so to complete the proof we may assume that both H_1 and H_2 are described as in item (i) or (ii) in Theorem 3.3. Write

$$\widehat{Q}(G, \tau) = \alpha + \beta,$$

where α is the contribution from involutions, and note that $|H_i| \leq 2\lceil \sqrt{n} \rceil!^2$ for $i = 1, 2$. Then by combining Lemmas 2.2 and 3.5 with Corollary 3.7, we obtain the bounds

$$\begin{aligned} \alpha &\leq i_2(H_1) i_2(H_2) \cdot \frac{2^{m/2} \lceil m/2 \rceil! (n - m)!}{n!} \\ &\leq \left(\lceil \sqrt{n} \rceil! (1 + 5^{(\sqrt{n}-1)/6})^2 \right)^2 \cdot \frac{2^{m/2} \lceil m/2 \rceil! (n - m)!}{n!} \end{aligned}$$

and

$$\beta \leq |H_1| |H_2| \cdot \frac{3^{m/3} \lceil m/3 \rceil! (n - m)!}{n!} \leq 4 \lceil \sqrt{n} \rceil!^4 \cdot \frac{3^{m/3} \lceil m/3 \rceil! (n - m)!}{n!},$$

where $m = \lceil 2\sqrt{n} \rceil$. Together, these bounds imply that $\alpha + \beta < 1$ and the result follows. \square

Part (iii) of Theorem 1 now follows as an easy corollary. Recall that $R_{\text{sol max}}(G)$ is the minimal integer k such that every k -tuple of soluble maximal subgroups of G is regular.

Proof of Theorem 1(iii). For $n \leq 16$ we can use MAGMA to verify the result (in Table 2 we record all the groups G with $R_{\text{sol max}}(G) = k > 2$, together with a non-regular soluble maximal $(k - 1)$ -tuple of the form (H, \dots, H)). And if $n \geq 17$, then every soluble maximal subgroup of G is primitive and therefore $R_{\text{sol max}}(G) = 2$ since $R_{\text{prim}}(G) = 2$ by part (ii) of Theorem 1. \square

$R_{\text{sol max}}(G)$	G	H
5	S_8	$S_4 \wr S_2$
4	S_5	S_4
	S_6	$S_3 \wr S_2$
3	$A_{6.2^2}$	$(S_3 \wr S_2).2$
	A_8	$(S_4 \wr S_2) \cap G$
	A_5	A_4
	A_6	$(S_3 \wr S_2) \cap G$
	$A_{6.2} = \text{PGL}_2(9)$	D_{20}
	$A_{6.2} = M_{10}$	$3^2:Q_8$
	A_7	$(S_4 \times S_3) \cap G$
	S_7	$S_4 \times S_3$
	A_9	$(S_3 \wr S_3) \cap G$
	S_9	$S_3 \wr S_3$
	A_{12}	$(S_4 \wr S_3) \cap G$
	S_{12}	$S_4 \wr S_3$
A_{16}	$(S_4 \wr S_4) \cap G$	
S_{16}	$S_4 \wr S_4$	

TABLE 2. Almost simple groups G with socle A_n and $R_{\text{sol max}}(G) > 2$

3.2. The regularity number. Let $G = S_n$ or A_n with $n \geq 5$. In this section we complete the proof of Theorem 1 by determining the regularity number of G . By considering the base size of the natural action of G on $[n] = \{1, \dots, n\}$, it suffices to show that $R(G) \leq k$, where $k = n - 1$ if $G = S_n$ and $k = n - 2$ if $G = A_n$. So in view of Lemma 2.9, we just need to verify that every k -tuple of subgroups in $\mathcal{M}(G)$ is regular, where $\mathcal{M}(G)$ is the set of core-free subgroups of G that are maximal in S_n or A_n .

The proof divides naturally into three cases since any subgroup in $\mathcal{M}(G)$ acts intransitively, imprimitively or primitively on $[n]$. Recall that we define $R_{\text{intrans}}(G)$ to be the minimal integer k such that every k -tuple of intransitive subgroups (H_1, \dots, H_k) of G is regular. Similarly, $R_{\text{imprim}}(G)$ is defined with respect to imprimitive subgroups (note that if n is a prime number, then every transitive subgroup of G is primitive, so $R_{\text{imprim}}(G)$ is not defined in this case).

In Section 3.2.1 we begin by handling the intransitive tuples, presenting several lemmas that will allow us to prove that $R_{\text{intrans}}(S_n) = n - 1$ and $R_{\text{intrans}}(A_n) = n - 2$ via a direct construction (see Proposition 3.10). The imprimitive tuples are then studied in Section 3.2.2. Here the analysis is more complicated and we rely on two key results (Lemmas 3.14 and 3.15), which may be of independent interest. Roughly speaking, our strategy is to reduce the imprimitive case to a situation involving intransitive subgroups, which we have already handled in Section 3.2.1. Finally, we complete the proof of Theorem 1 by combining our work on intransitive and imprimitive tuples with our results on primitive subgroups in Section 3.1. The details are presented in Section 3.2.3.

3.2.1. Intransitive tuples. Our main result in this section is Proposition 3.10, which gives

$$R_{\text{intrans}}(S_n) = n - 1 \quad \text{and} \quad R_{\text{intrans}}(A_n) = n - 2.$$

Throughout this section we fix an integer $n \geq 5$ and we define W_n to be the set of subsets of $[n] = \{1, \dots, n\}$ of size at most $n/2$. Following Halasi [31], if $\alpha \in [n]$ and $\mathcal{A} \subseteq W_n$, then we refer to

$$N_{\mathcal{A}}(\alpha) = \{X \in \mathcal{A} : \alpha \in X\}$$

as the *neighbourhood* of α with respect to \mathcal{A} .

Lemma 3.8. *Let $\mathcal{A} = \{X_1, \dots, X_k\} \subseteq W_n$ and set $H = \bigcap_{i=1}^k H_i$, where H_i is the setwise stabiliser of X_i in S_n .*

- (i) *For all $\alpha, \beta \in [n]$, we have $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$ if and only if α and β are contained in different H -orbits.*

(ii) *In particular, $H = 1$ if and only if $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$ for all distinct $\alpha, \beta \in [n]$.*

Proof. Since (ii) follows immediately from (i), we just need to establish the claim in (i). First assume α and β lie in distinct orbits of H . Then H does not contain the transposition $g = (\alpha, \beta) \in S_n$, which means that $X_i^g \neq X_i$ for some i . Therefore, either $X_i \in N_{\mathcal{A}}(\alpha) \setminus N_{\mathcal{A}}(\beta)$ or $X_i \in N_{\mathcal{A}}(\beta) \setminus N_{\mathcal{A}}(\alpha)$, and thus $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$.

Conversely, suppose $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$. Then we may assume there exists a set $X_i \in \mathcal{A}$ such that $\alpha \in X_i$ and $\beta \notin X_i$. Consider an element $g \in S_n$ such that $\alpha^g = \beta$. Then $X_i^g \neq X_i$ and thus $g \notin H$, which in turn implies that α and β are in distinct H -orbits. \square

We now present a lemma that will play a key role in the proof of Proposition 3.10. It will also be used repeatedly in the proof of Theorem 1 in Section 3.2.3.

Lemma 3.9. *Let a_1, \dots, a_k be positive integers such that $a_1 \leq a_2 \leq \dots \leq a_k \leq n/2$, where $2 \leq k < n$. For each $i \in \{1, \dots, k\}$, let X_i be the following a_i -element subset of $[n]$ and let $H = \bigcap_{i=1}^k H_i$, where H_i is the stabiliser of X_i in S_n :*

$$\begin{aligned} X_1 &= \{1, \dots, a_1\} \\ X_2 &= \{2, \dots, a_2 + 1\} \\ &\vdots \\ X_{k-1} &= \{k-1, \dots, n-1, 1, 2, \dots, a_{k-1} - (n-k+1)\} \\ X_k &= \{k, \dots, n-1, 1, 2, \dots, a_k - (n-k)\} \end{aligned} \tag{1}$$

Then $H \leq \text{Sym}(\{k, \dots, m\}) \times \text{Sym}(\{m+1, \dots, n\})$, where $m = \max\{\alpha : \alpha \in X_k\}$.

Proof. Let $\mathcal{A} = \{X_1, \dots, X_k\}$. First we show that $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$ for all distinct $\alpha, \beta \in \{1, \dots, k, n\}$. To do this, we may assume $\alpha < \beta$.

If $\beta = n$, then $N_{\mathcal{A}}(\beta)$ is empty, whereas $X_\alpha \in N_{\mathcal{A}}(\alpha)$, whence $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$. Now assume $\beta \leq k$ and suppose that $\alpha \in X_\beta$ and $\beta \in X_\alpha$. The containment of α in X_β implies that $(n-1) - \beta + 1 + \alpha \leq a_\beta$ and thus $\beta - \alpha \geq n/2$ since $a_\beta \leq n/2$. Similarly, since $\beta \in X_\alpha$ we have $\beta - \alpha + 1 \leq a_\alpha$, so $\beta - \alpha < n/2$ and we have reached a contradiction. It follows that either $\alpha \notin X_\beta$ or $\beta \notin X_\alpha$, whence $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$ as required.

With a very similar argument, we can show that $N_{\mathcal{A}}(\alpha) \neq N_{\mathcal{A}}(\beta)$ for all $\alpha \in \{1, \dots, k-1\}$ and $\beta \in \{k, \dots, m\}$. In addition, we claim that $N_{\mathcal{A}}(\gamma)$ is empty for all $\gamma \geq m+1$. This is clear if $m = n-1$, so we may assume $m < n-1$ and thus $X_k = \{k, k+1, \dots, m\}$. For each $i \in \{1, \dots, k\}$ we have

$$|X_i| = a_i \leq a_k = m - k + 1 < \gamma - k + 1,$$

so $\gamma \notin X_i$ and $N_{\mathcal{A}}(\gamma)$ is indeed empty as claimed. In particular, since $i \in X_i$ and $\delta \in X_k$ for all $\delta \in \{k, \dots, m\}$, this means that $N_{\mathcal{A}}(\gamma) \neq N_{\mathcal{A}}(\delta)$ for all $\gamma \geq m+1$ and all $\delta \in \{1, \dots, m\}$.

Putting all of this together and applying Lemma 3.8, we deduce the following:

- (a) The points $1, \dots, k, n$ are contained in distinct H -orbits.
- (b) Each $\alpha \in \{1, \dots, k-1\}$ and $\beta \in \{k, \dots, m\}$ are contained in distinct H -orbits.
- (c) Each $\alpha \in \{1, \dots, m\}$ and $\beta \in \{m+1, \dots, n\}$ are contained in distinct H -orbits.

As a consequence, H acts trivially on $\{1, \dots, k-1\}$ and it preserves the subsets $\{k, \dots, m\}$ and $\{m+1, \dots, n\}$. In other words,

$$H \leq \text{Sym}(\{k, \dots, m\}) \times \text{Sym}(\{m+1, \dots, n\})$$

as required. \square

We are now in a position to establish our main result on intransitive tuples.

Proposition 3.10. *For all $n \geq 5$, we have $R_{\text{intrans}}(S_n) = n-1$ and $R_{\text{intrans}}(A_n) = n-2$.*

Proof. First observe that the intransitive $(n-2)$ -tuple $(S_{n-1}, \dots, S_{n-1})$ is non-regular, so $R_{\text{intrans}}(S_n) \geq n-1$. Similarly, $R_{\text{intrans}}(A_n) \geq n-2$, so we just need to establish the upper bounds $R_{\text{intrans}}(S_n) \leq n-1$ and $R_{\text{intrans}}(A_n) \leq n-2$.

Suppose $G = S_n$ or A_n , and let $\tau = (K_1, \dots, K_{n-2})$ be an $(n-2)$ -tuple of intransitive subgroups of G . We may embed each K_i in the setwise stabiliser $L_i = S_{a_i} \times S_{n-a_i}$ in S_n of some subset of $[n]$ of size $a_i \leq n/2$, and by reordering we may assume that $a_i \leq a_{i+1}$ for all i . We claim that there exist elements $g_i \in G$ such that

$$\bigcap_{i=1}^{n-2} L_i^{g_i} \leq \langle (\alpha, \beta) \rangle, \quad (2)$$

where $\alpha = n-1$ and $\beta \in \{n-2, n\}$. Since G acts transitively on the set of k -element subsets of $[n]$ for any fixed $k \leq n/2$, we just need to define a collection of subsets X_1, \dots, X_{n-2} of $[n]$ such that $|X_i| = a_i$ and $H = \bigcap_{i=1}^{n-2} H_i \leq \langle (\alpha, \beta) \rangle$, where H_i is the setwise stabiliser of X_i in S_n . To do this, we can define the sets X_1, \dots, X_{n-2} as in (1) (with $k = n-2$), and then by applying Lemma 3.9 we deduce that $H \leq \langle (\alpha, \beta) \rangle$ as required.

For $G = A_n$ we immediately deduce that τ is regular and thus $R_{\text{intrans}}(A_n) \leq n-2$. So for the remainder, let us assume $G = S_n$ and let $\tau' = (K_1, \dots, K_{n-1})$ be an intransitive $(n-1)$ -tuple. By the above argument, we have $\bigcap_{i=1}^{n-2} K_i^{g_i} \leq \langle (\alpha, \beta) \rangle$ for some $g_i \in G$. Now K_{n-1} is contained in the setwise stabiliser in G of some k -element subset of $[n]$ with $k \leq n/2$. Of course, we can choose a k -set X_{n-1} containing α but not β , in which case the stabiliser of X_{n-1} in G does not contain the transposition (α, β) . In other words, there exists an element $g_{n-1} \in G$ such that

$$\bigcap_{i=1}^{n-1} K_i^{g_i} \leq \langle (\alpha, \beta) \rangle \cap K_{n-1}^{g_{n-1}} = 1$$

and we conclude that τ' is regular. This completes the proof of the proposition. \square

3.2.2. Imprimitive tuples. In this section we focus on imprimitive tuples and our main result is Proposition 3.18, which states that

$$R_{\text{imprim}}(S_n) \leq n-1 \quad \text{and} \quad R_{\text{imprim}}(A_n) \leq n-2$$

for all composite integers $n \geq 6$ (recall that if n is a prime, then every transitive subgroup of S_n is primitive). To establish this bound for $G = S_n$, we need to prove that every $(n-1)$ -tuple of the form $(S_{a_1} \wr S_{b_1}, \dots, S_{a_{n-1}} \wr S_{b_{n-1}})$ is regular, where $n = a_i b_i$ and $a_i \geq 2$ for all i . And similarly for $G = A_n$ with respect to $(n-2)$ -tuples with components of the form $(S_{a_i} \wr S_{b_i}) \cap G$. Throughout this section we assume $n \geq 6$ is composite and we write $[n] = \{1, \dots, n\}$.

Remark 3.11. Let $G = S_n$ or A_n with $n \geq 6$ composite. We have not attempted to seek sharper estimates on $R_{\text{imprim}}(G)$ because the above bounds (together with some of the auxiliary results presented below) will be sufficient for our proof of Theorem 1(i). However, it seems reasonable to expect that the bounds in Proposition 3.18 are far from best possible. Indeed, the main theorem of [43] yields

$$B_{\text{imprim}}(G) \leq \log_2 n + 2,$$

which leads us to speculate that $R_{\text{imprim}}(G) = O(\log_2 n)$. With the aid of MAGMA, we have checked that $R_{\text{imprim}}(G) \leq 6$ for all $n < 32$, but here we do not pursue this any further.

We begin by defining some useful terminology that arises naturally in this setting. We will say that a partition X of $[n]$ is *uniform* if every part has size ℓ , where $1 < \ell < n$, and we will refer to X as an ℓ -*partition*. If X is an ℓ -partition, then the stabiliser of X in S_n is defined to be the largest subgroup $H \leq S_n$ that preserves the partition, so we have $H = S_\ell \wr S_{n/\ell}$. Similarly, the stabiliser in A_n is the subgroup $H \cap A_n$.

Let X_1 and X_2 be uniform partitions of $[n]$ with parts of size a and b , respectively. Choose an ordering of the parts P_1, \dots, P_k comprising X_1 , where $k = n/a$, and assign the colour c_i

to every point in P_i , where c_1, \dots, c_k are distinct. Then the corresponding *colour function* is the map $\chi : [n] \rightarrow \{c_1, \dots, c_k\}$. We say that a part Q of X_2 has *colour sequence* $(c_{q_1}, \dots, c_{q_b})$ if there is a bijection from Q to the multiset of colours $\{c_{q_1}, \dots, c_{q_b}\}$ such that each point in Q is mapped to its assigned colour. And if R is another part of X_2 with colour sequence $(c_{r_1}, \dots, c_{r_b})$, we will say that Q and R are *colour-equivalent* if there exists a permutation of the colour set $\{c_1, \dots, c_k\}$ mapping the multiset $\{c_{q_1}, \dots, c_{q_b}\}$ to the multiset $\{c_{r_1}, \dots, c_{r_b}\}$. Finally, we say that a part of X_2 is *monochrome* if all of its points have the same colour.

We begin with an easy lemma.

Lemma 3.12. *Let X_1, X_2 be defined as above and let H_i be the stabiliser of X_i in S_n . If $g \in H_1 \cap H_2$ and Q is a part of X_2 , then Q and Q^g are colour-equivalent.*

Proof. Since $g \in H_1$, it follows that any two elements in $[n]$ with the same colour are mapped under g to elements of the same colour. In particular, Q and Q^g are colour-equivalent. \square

Lemma 3.13. *Let X be an ℓ -partition of $[n]$ with parts P_1, \dots, P_k , where $\ell \geq 3$. Fix $\alpha \in P_i$ and $\beta \in P_j$, where $i \neq j$, and let $Y = X^t$ be the ℓ -partition with parts $Q_i = P_i^t$, where $t = (\alpha, \beta) \in S_n$. If H_X and H_Y denote the stabilisers of X and Y in S_n , then $H_X \cap H_Y$ fixes $\{\alpha, \beta\}$ setwise.*

Proof. For each $s \in \{1, \dots, k\}$, assign the colour c_s to the points in P_s , where c_1, \dots, c_k are distinct. Note that

$$Q_s = \begin{cases} P_s & \text{if } s \neq i, j \\ (P_i \setminus \{\alpha\}) \cup \{\beta\} & \text{if } s = i \\ (P_j \setminus \{\beta\}) \cup \{\alpha\} & \text{if } s = j \end{cases}$$

for all $s \in \{1, \dots, k\}$. Fix an element $g \in H_X \cap H_Y$.

Suppose $Q_i^g \neq Q_i$. Then since $Q_s = P_s$ is monochrome for all $s \neq i, j$, Lemma 3.12 implies that $Q_i^g = Q_j$ is the only option. Moreover, since any two points in $[n]$ of the same colour must be mapped to points of the same colour by any element in H_X , it follows that $\beta^g = \alpha$. Similarly, if $Q_i^g = Q_i$ then $\beta^g = \beta$.

If we now repeat the above argument, with Q_i replaced by Q_j , we deduce that either g interchanges Q_i and Q_j , in which case $\beta^g = \alpha$ and $\alpha^g = \beta$, or g fixes both Q_i and Q_j , which means that $\alpha^g = \alpha$ and $\beta^g = \beta$. The result follows. \square

As an application of Lemma 3.13 we can now establish the following technical result, which will be an important ingredient in our proof of Proposition 3.18.

Lemma 3.14. *Suppose $n = 2\ell$ is even and A is an m -element subset of $[n]$ with $3 \leq m < n$. Then there exist ℓ -partitions X_1, \dots, X_m of $[n]$ such that $\bigcap_{i=1}^m H_i$ fixes A pointwise, where H_i is the stabiliser of X_i in S_n .*

Proof. It suffices to show that there exists a collection of ℓ -partitions Y_1, \dots, Y_m with respective stabilisers L_1, \dots, L_m in S_n such that $L = \bigcap_{i=1}^m L_i$ fixes some m -set B pointwise. Indeed, the partitions $X_i = Y_i^g$ will then have the required property, where $g \in S_n$ is chosen such that $B^g = A$.

We will define appropriate partitions Y_i , with parts labelled $P_{i,1}$ and $P_{i,2}$, and stabiliser L_i in S_n . For Y_1 and Y_2 we set

$$P_{1,1} = \{1, 2, \dots, \ell\}, \quad P_{1,2} = \{\ell + 1, \ell + 2, \dots, n\}$$

and

$$P_{2,1} = \{2, 3, \dots, \ell + 1\}, \quad P_{2,2} = \{1, \ell + 2, \dots, n\}.$$

We then define Y_3, \dots, Y_m iteratively, treating the cases $m \leq \ell + 1$ and $m > \ell + 1$ separately.

First assume $m \leq \ell + 1$. Here we define

$$P_{i,1} = \{1, \dots, \ell + 1\} \setminus \{i - 1\}, \quad P_{i,2} = \{\ell + 2, \dots, n\} \cup \{i - 1\} \quad (3)$$

for $i \geq 3$, which means that

$$P_{2,1} = (P_{1,1} \setminus \{1\}) \cup \{\ell + 1\}, \quad P_{2,2} = (P_{1,2} \setminus \{\ell + 1\}) \cup \{1\}$$

and

$$P_{i,1} = (P_{i-1,1} \setminus \{i-1\}) \cup \{i-2\}, \quad P_{i,2} = (P_{i-1,2} \setminus \{i-2\}) \cup \{i-1\}$$

for all $i \geq 3$. Therefore, Lemma 3.13 implies that $L_1 \cap L_2$ fixes $\{1, \ell+1\}$ setwise, and similarly $L_{i-1} \cap L_i$ fixes $\{i-2, i-1\}$ setwise for each $i \geq 3$. In particular, $L = \bigcap_{i=1}^m L_i$ fixes each of the following 2-sets

$$\{1, \ell+1\}, \{1, 2\}, \{2, 3\}, \dots, \{m-2, m-1\},$$

and hence L fixes the m -set $\{1, 2, \dots, m-1, \ell+1\}$ pointwise.

Finally, suppose $m > \ell+1$. For $3 \leq i \leq \ell+1$ we define the two parts of Y_i as in (3), and for $i > \ell+1$ we set

$$P_{i,1} = \{2, \dots, \ell\} \cup \{i\}, \quad P_{i,2} = \{1, \ell+1, \dots, n\} \setminus \{i\}.$$

As above, we deduce that $\bigcap_{i=1}^{\ell+1} L_i$ fixes $\{1, \dots, \ell+1\}$ pointwise. For $i = \ell+2$, note that

$$P_{i,1} = (P_{2,1} \setminus \{i-1\}) \cup \{i\}, \quad P_{i,2} = (P_{2,2} \setminus \{i\}) \cup \{i-1\}$$

and for $i > \ell+2$ we have

$$P_{i,1} = (P_{i-1,1} \setminus \{i-1\}) \cup \{i\}, \quad P_{i,2} = (P_{i-1,2} \setminus \{i\}) \cup \{i-1\}.$$

Therefore, by applying Lemma 3.13, we deduce that $\bigcap_{i=\ell+2}^m L_i$ fixes each of the 2-sets

$$\{\ell+1, \ell+2\}, \{\ell+2, \ell+3\}, \dots, \{m-1, m\}$$

setwise and thus $L = \bigcap_{i=1}^m L_i$ fixes $\{1, \dots, m\}$ pointwise. \square

We are now ready to present the following lemma, which will play an essential role in our proof of Proposition 3.18. It will also be used in Section 3.2.3, where we complete the proof of Theorem 1. The proof is long and technical, so we partition the argument into a number of separate and clearly defined cases.

Lemma 3.15. *Let $G = S_n$ or A_n , and let X_i be an a_i -partition of $[n]$ for $i = 1, 2$, where $a_1 \geq a_2$. Let $\alpha, \beta \in [n]$ be distinct and assume that either*

- (i) $n/2 > a_1 \geq 3$; or
- (ii) $a_1 = n/2$, $a_2 \geq 3$ and α and β are contained in different parts of X_1 .

Then there exists an element $g \in G$ such that $H_1 \cap H_2^g$ fixes $\{\alpha, \beta\}$ setwise, where H_i is the stabiliser of X_i in G .

Proof. Write $k = n/a_1$, $m = n/a_2$ and let P_1, \dots, P_k be the parts of X_1 . We assign the colour c_i to the points in P_i , where c_1, \dots, c_k are distinct, and we let χ be the corresponding colour function on $[n]$. In order to prove the lemma, we need to construct an a_2 -partition X_3 of $[n]$ with stabiliser H_3 such that $H_1 \cap H_3$ fixes $\{\alpha, \beta\}$ setwise. Indeed, the transitivity of G on the set of a_2 -partitions of $[n]$ implies that $H_3 = H_2^g$ for some $g \in G$ and the desired result follows.

First assume (i) holds, so $n/2 > a_1 \geq 3$ and $k = n/a_1 \geq 3$. There are two different cases to consider, according to whether or not $a_2 = 2$.

Case 1. $n/2 > a_1 \geq a_2 \geq 3$

We begin by assuming $a_2 \geq 3$. We will write $P_i = \{p_{i1}, \dots, p_{ia_1}\}$ for each i and we consider two subcases, according to the position of α and β in the parts of X_1 .

Case 1(a). α and β are in different parts of X_1

Without loss of generality, we may assume that $\alpha = p_{11} \in P_1$ and $\beta = p_{21} \in P_2$. We first construct an a_2 -partition Y of $[n]$ with parts Q_1, \dots, Q_m and stabiliser H_Y in G by setting

$$Q_1 = \{\alpha, p_{22}, \dots, p_{2a_2}\}, \quad Q_2 = \{\beta, p_{12}, \dots, p_{1a_2}\}$$

and we define Q_3, \dots, Q_m by partitioning the remaining points in $[n]$ into sets of size a_2 in the following order:

$$P_{1(a_2+1)}, \dots, P_{1a_1}, P_{31}, \dots, P_{3a_1}, P_{2(a_2+1)}, \dots, P_{2a_1}, P_{41}, \dots, P_{4a_1}, \dots, P_{k1}, \dots, P_{ka_1}.$$

Notice that Q_1 and Q_2 are the only parts of Y with respective colour sequences (c_1, c_2, \dots, c_2) and (c_2, c_1, \dots, c_1) . Moreover, if $i \neq j$ then our construction implies that Y has parts with colour sequences (c_i, c_j, \dots, c_j) and (c_j, c_i, \dots, c_i) if and only if $\{i, j\} = \{1, 2\}$.

Let $x \in H_1 \cap H_Y$. Then Lemma 3.12 implies that Q_1^x has colour sequence (c_i, c_j, \dots, c_j) for some $i \neq j$. Since $a_2 \geq 3$, we deduce that $P_1^x = P_i$ and $P_2^x = P_j$, which in turn implies that Q_2^x has colour sequence (c_j, c_i, \dots, c_i) . So as noted above, we have $(i, j) = (1, 2)$ or $(2, 1)$. If $(i, j) = (1, 2)$ then $Q_1^x = Q_1$ and we deduce that $\alpha^x = \alpha$. On the other hand, if $(i, j) = (2, 1)$ then $Q_1^x = Q_2$ and $\alpha^x = \beta$.

Similarly, by repeating the argument with Q_1 replaced by Q_2 , we deduce that $\beta^x \in \{\alpha, \beta\}$. Therefore, $H_1 \cap H_Y$ fixes $\{\alpha, \beta\}$ setwise, as required.

Case 1(b). α and β are contained in the same part of X_1

By relabelling, we may assume that $\alpha, \beta \in P_1$, say $\alpha = p_{11}$ and $\beta = p_{12}$. We now define a new a_2 -partition Y of $[n]$ with parts Q_1, \dots, Q_m and stabiliser H_Y in G . To construct Y , we first set

$$Q_1 = \{\alpha, p_{21}, \dots, p_{2(a_2-1)}\}, \quad Q_2 = \{\beta, p_{31}, \dots, p_{3(a_2-1)}\},$$

so Q_1 and Q_2 have respective colour sequences (c_1, c_2, \dots, c_2) and (c_1, c_3, \dots, c_3) . We then divide the remaining points of $[n]$ into parts Q_3, \dots, Q_m of size a_2 in such a way that none of these parts has colour sequence (c_1, c_i, \dots, c_i) with $i \geq 2$.

Observe that we can always construct such a partition. To justify this claim, first assume $a_1 = 3$, so $a_2 = 3$ and $k = m \geq 3$. Here we can set $Q_3 = \{p_{13}, p_{23}, p_{33}\}$ and then define Q_4, \dots, Q_m arbitrarily, noting that none of the latter parts will contain any points with colour c_1 . Now assume $a_1 \geq 4$ and write $a_1 - 2 = sa_2 + r$, where $s \geq 0$ and $0 \leq r < a_2$. Here we can define Q_3, \dots, Q_{s+2} so that they all have colour sequence (c_1, \dots, c_1) . If $r \neq 1$ then we can define Q_{s+3} so that it contains exactly r points with colour c_1 , while the remaining parts Q_{s+4}, \dots, Q_m contain no points from P_1 . And if $r = 1$ then we can choose Q_{s+3} so that it contains points from P_1, P_2 and P_3 , while none of the remaining parts in Y contain any points with colour c_1 . This justifies the claim.

Before continuing the analysis of Case 1(b), let us pause to provide a brief outline of the argument. We are seeking to construct an a_2 -partition W of $[n]$ with stabiliser H_W , containing the parts Q_1 and Q_2 defined above, with the property that $\{Q_1, Q_2\}^x = \{Q_1, Q_2\}$ for all $x \in H_1 \cap H_W$. Indeed, if we have such a partition, then $H_1 \cap H_W$ fixes $\{\alpha, \beta\}$ and the result follows. Set $Y_0 = Y$, which is the a_2 -partition defined above. If Y_0 does not have the desired property, then we will construct a new partition $Y_1 = Y_0^g$ with parts Q_i^g for some carefully chosen transposition $g \in S_n$. We can then repeat this process and we will prove that it produces a partition $W = Y_t$ with the desired property after finitely many iterations.

We start by inspecting the partition Y constructed above. Fix an element $x \in H_1 \cap H_Y$ and consider the image of Q_1 under x .

First assume $Q_1^x = Q_1$. Here $P_1^x = P_1$ and $P_2^x = P_2$ since $a_2 \geq 3$, whence $\alpha^x = \alpha$ and Q_2^x has colour sequence (c_1, c_i, \dots, c_i) for some $i \geq 2$. Then by the construction of Y , it follows that $Q_2^x = Q_2$ and thus $\beta^x = \beta$. Similarly, if $Q_1^x = Q_2$ then $P_1^x = P_1$ and $\alpha^x = \beta$, which in turn implies that $Q_2^x = Q_1$ and $\beta^x = \alpha$.

So if $Q_1^x \in \{Q_1, Q_2\}$ for all $x \in H_1 \cap H_Y$, then $H_1 \cap H_Y$ fixes $\{\alpha, \beta\}$ and we are done. So let us assume there exists an element $x \in H_1 \cap H_Y$ such that $Q_1^x = Q_a$ for some $a \geq 3$. Now Q_1^x has colour sequence (c_i, c_j, \dots, c_j) , where $i \neq j$ and $i \geq 2$. More precisely, $\chi(\alpha^x) = c_i$ and therefore $\chi(\beta^x) = c_i$ since α and β are in the same part of X_1 . It follows that Q_2^x has colour sequence $(c_i, c_\ell, \dots, c_\ell)$ for some $\ell \neq i$, and we note that $\ell \neq j$ since $P_2^x \neq P_3^x$. Fix $\gamma_1 \in Q_1^x$ and $\delta_1 \in Q_2^x$ such that $\chi(\gamma_1) = c_j$ and $\chi(\delta_1) = c_\ell$, and let $g_1 \in S_n$ be the transposition (γ_1, δ_1) . Note that the parts of Y containing γ_1 and δ_1 are both colour-equivalent to Q_1 .

At this point, we now switch to the a_2 -partition $Z = Y_1 = Y^{g_1}$, with parts $R_\ell = Q_\ell^{g_1}$ for $1 \leq \ell \leq m$. Let H_Z be the stabiliser of Z in G and let $x \in H_1 \cap H_Z$. Notice that $R_1 = Q_1$ and $R_2 = Q_2$ since $\gamma_1, \delta_1 \notin Q_1 \cup Q_2$.

By arguing as above, we deduce that either $R_1^x = R_1$ and $\alpha^x = \alpha$, or $R_1^x = R_2$ and $\alpha^x = \beta$, or $R_1^x = R_a$ for some $a \geq 3$. Let us assume that we are in the latter situation. Then as above, we note that R_1^x has colour sequence (c_i, c_j, \dots, c_j) for some $i \geq 2$ with $i \neq j$, and it follows that R_2^x has colour sequence $(c_i, c_\ell, \dots, c_\ell)$ for some $\ell \neq i, j$. We now choose $\gamma_2 \in R_1^x$ and $\delta_2 \in R_2^x$ such that $\chi(\gamma_2) = c_j$ and $\chi(\delta_2) = c_\ell$, and we construct the a_2 -partition $Y_2 = Y_1^{g_2} = Y^{g_1 g_2}$, where $g_2 = (\gamma_2, \delta_2) \in S_n$.

We can now repeat this process until after t steps we obtain an a_2 -partition $W = Y_t = Y^g$ of $[n]$ containing the parts Q_1 and Q_2 , where $g = g_1 \cdots g_t$ is a product of transpositions and $Q_1^x \in \{Q_1, Q_2\}$ for all $x \in H_1 \cap H_W$. We claim that this process does indeed produce such a partition after finitely many iterations. To see this, first observe that if $a \in \{2, \dots, t\}$, then the parts comprising the partition $Y_a = Y^{g_1 \cdots g_a}$ are as follows:

$$Q_j^{g_1 \cdots g_a} = \begin{cases} Q_j^{g_1 \cdots g_{a-1}} & \text{if } \{\gamma_a, \delta_a\} \notin Q_j^{g_1 \cdots g_{a-1}} \\ Q' = (Q_j^{g_1 \cdots g_{a-1}} \setminus \{\gamma_a\}) \cup \{\delta_a\} & \text{if } \gamma_a \in Q_j^{g_1 \cdots g_{a-1}} \\ Q'' = (Q_j^{g_1 \cdots g_{a-1}} \setminus \{\delta_a\}) \cup \{\gamma_a\} & \text{if } \delta_a \in Q_j^{g_1 \cdots g_{a-1}} \end{cases} \quad (4)$$

In particular, notice that the parts of $Y_{a-1} = Y^{g_1 \cdots g_{a-1}}$ containing γ_a and δ_a are both colour-equivalent to $Q_1 = Q_1^{g_1 \cdots g_{a-1}}$, whereas Q' and Q'' are not. Therefore, the number of parts that are colour-equivalent to Q_1 strictly decreases with each iteration, and this justifies the claim since our initial partition Y has only finitely many parts.

So we now have an a_2 -partition W containing the parts Q_1 and Q_2 such that $Q_1^x \in \{Q_1, Q_2\}$ for all $x \in H_1 \cap H_W$. In particular, this implies that $P_1^x = P_1$ and $\alpha^x \in \{\alpha, \beta\}$ for all $x \in H_1 \cap H_W$. So to complete the analysis of Case 1(b), it suffices to show that Q_1 and Q_2 are the only parts in W with colour sequence (c_1, c_i, \dots, c_i) for some $i \geq 2$. Indeed, if this property is satisfied then $Q_2^x \in \{Q_1, Q_2\}$ and $\beta^x \in \{\alpha, \beta\}$ for all $x \in H_1 \cap H_W$, which means that every element in $H_1 \cap H_W$ fixes $\{\alpha, \beta\}$ setwise, as required.

So it just remains to justify the colour sequence claim, which we will do via induction on t , recalling that $W = Y_t = Y^g$ and $g = g_1 \cdots g_t$ as defined above. For the base case $t = 0$ we have $W = Y$ and we know that Y has the desired property by construction. Now assume $t \geq 1$ and suppose that Q_1 and Q_2 are the only parts in Y_{t-1} with colour sequence (c_1, c_i, \dots, c_i) for some $i \geq 2$. From (4), we see that every part in Y_t , other than Q' and Q'' , is also a part of Y_{t-1} , so Q_1 and Q_2 are the only parts in $Y_t \setminus \{Q', Q''\}$ with colour sequence (c_1, c_i, \dots, c_i) for some $i \geq 2$. The claim now follows since it is clear to see that neither Q' nor Q'' is colour-equivalent to Q_1 or Q_2 .

This completes the proof of the lemma in Case 1.

Case 2. $n/2 > a_1 \geq 3$, $a_2 = 2$

In order to complete the proof of the lemma in case (i), we may assume $a_2 = 2$. As above, set $k = n/a_1$ and let P_1, \dots, P_k be the parts of X_1 . For each i , it will be convenient to write $p_{(i-1)a_1+1}, \dots, p_{ia_1}$ for the points in P_i , to which we assign the colour c_i as before. Let $\chi : [n] \rightarrow \{c_1, \dots, c_k\}$ be the corresponding colour function.

Case 2(a). α and β are contained in the same part of X_1

We may assume $\alpha, \beta \in P_1$, say $\alpha = p_1$ and $\beta = p_2$. Consider the 2-partition Y of $[n]$ with parts

$$\{p_1, p_2\}, \{p_3, p_{n/2+2}\}, \{p_4, p_{n/2+3}\}, \dots, \{p_{n/2+1}, p_n\}$$

and let H_Y be the stabiliser of Y in G . Note that every part of Y , other than $\{p_1, p_2\}$, is of the form $\{p_i, p_{n/2-1+i}\}$ for some $i \geq 3$. Since the parts in X_1 have size $a_1 < n/2$, it follows that p_i and $p_{n/2-1+i}$ cannot both be contained in the same part of X_1 . Therefore, no part

of Y , other than $\{p_1, p_2\}$, is colour-equivalent to $\{\alpha, \beta\}$, and by applying Lemma 3.12 we deduce that $\{\alpha, \beta\}$ is fixed by $H_1 \cap H_Y$.

Case 2(b). α and β are contained in different parts of X_1

We may assume $\alpha \in P_1$ and $\beta \in P_2$, say $\alpha = p_1$ and $\beta = p_{a_1+1}$. We now consider two subcases, according to the parity of a_1 .

First assume $a_1 \geq 3$ is odd, which means that $k = n/a_1$ is even. Define a 2-partition Y of $[n]$ with the following parts

$$\{p_1, p_{a_1+1}\}, \{p_2, p_3\}, \dots, \{p_{a_1-1}, p_{a_1}\}, \{p_{a_1+2}, p_{a_1+3}\}, \dots, \{p_{2a_1-1}, p_{2a_1}\}, \\ \{p_{2a_1+1}, p_{3a_1+1}\}, \dots, \{p_{3a_1}, p_{4a_1}\}, \dots, \{p_{n-2a_1+1}, p_{n-a_1+1}\}, \dots, \{p_{n-a_1}, p_n\}$$

Note that every part in the second row is of the form $\{p_i, p_{a_1+i}\}$ for some $i \geq 2a_1 + 1$. We claim that each $x \in H_1 \cap H_Y$ fixes $\{\alpha, \beta\}$ setwise.

First note that $\{\alpha, \beta\}$ cannot be mapped by x to any of the following parts

$$\{p_2, p_3\}, \dots, \{p_{a_1-1}, p_{a_1}\}, \{p_{a_1+2}, p_{a_1+3}\}, \dots, \{p_{2a_1-1}, p_{2a_1}\}$$

since the latter are all monochrome (see Lemma 3.12). Now assume $\{\alpha, \beta\}^x = \{p_i, p_{a_1+i}\}$ for some $i \geq 2a_1 + 1$. Then since $\alpha \in P_1$ and $\chi(\alpha^x) \neq c_1, c_2$, it follows that $P_1^x = P_j$ for some $j \geq 3$ and thus $\{p_2, p_3\} \subseteq P_1$ must be mapped by x to some part of Y with colour sequence (c_j, c_j) . But no part of Y has such a colour sequence, so this possibility cannot arise and we conclude that $H_1 \cap H_Y$ fixes $\{\alpha, \beta\}$ setwise.

To complete the argument in Case 2(b), we may assume $a_1 \geq 4$ is even. Here we define a 2-partition Y with parts

$$\{p_1, p_{2a_1+1}\}, \{p_{a_1+1}, p_{2a_1+2}\}, \{p_2, p_{a_1+2}\}, \dots, \{p_{a_1}, p_{2a_1}\}, \{p_{2a_1+3}, p_{2a_1+4}\}, \dots, \{p_{n-1}, p_n\}$$

and our goal is to show that $H_1 \cap H_Y$ fixes $\{\alpha, \beta\}$ setwise. Let $x \in H_1 \cap H_Y$ and recall that $\alpha = p_1$ and $\beta = p_{a_1+1}$. We proceed by considering the images of the parts $\{p_1, p_{2a_1+1}\}$ and $\{p_{a_1+1}, p_{2a_1+2}\}$ under x .

First observe that $\{p_1, p_{2a_1+1}\}$ and $\{p_{a_1+1}, p_{2a_1+2}\}$ are non-monochrome, so neither of them can be moved by x to any of the monochrome parts $\{p_{2a_1+3}, p_{2a_1+4}\}, \dots, \{p_{n-1}, p_n\}$.

Next assume $\{p_1, p_{2a_1+1}\}^x = \{p_i, p_{a_1+i}\}$ for some $i \in \{2, \dots, a_1\}$. This implies that the point $p_{2a_1+1} \in P_3$ has colour c_1 or c_2 , so $P_3^x \in \{P_1, P_2\}$ and thus $\{p_{2a_1+3}, p_{2a_1+4}\}^x$ has colour sequence (c_j, c_j) for some $j \in \{1, 2\}$. However, there is no such part in Y , so this situation does not arise. This means that $\{p_1, p_{2a_1+1}\}^x \neq \{p_i, p_{a_1+i}\}$ with $i \in \{2, \dots, a_1\}$, and similarly we deduce that the same conclusion holds for $\{p_{a_1+1}, p_{2a_1+2}\}^x$.

We have now shown that x either fixes the parts $\{p_1, p_{2a_1+1}\}$ and $\{p_{a_1+1}, p_{2a_1+2}\}$, or it interchanges them.

We claim that if x fixes $\{p_1, p_{2a_1+1}\}$, then it must fix $p_1 = \alpha$. Suppose otherwise, so $\alpha^x = p_{2a_1+1}$. Since $x \in H_1$, it follows that $P_1^x = P_3$, so the parts $\{p_2, p_{a_1+2}\}, \dots, \{p_{a_1}, p_{2a_1}\}$ must all be mapped by x to some part of Y with colour sequence (c_3, c_i) for some $i \neq 3$. However, there is only one such part, namely $\{p_{a_1+1}, p_{2a_1+2}\}$, so we have reached a contradiction since $a_1 \geq 3$. Similarly, since x fixes $\{p_{a_1+1}, p_{2a_1+2}\}$, we deduce that $\beta^x = \beta$.

Finally, let us assume x interchanges the parts $\{p_1, p_{2a_1+1}\}$ and $\{p_{a_1+1}, p_{2a_1+2}\}$. As in the previous paragraph, we know that $P_1^x \neq P_3$, so we must have $\alpha^x = p_1^x = p_{a_1+1} = \beta$. This implies that $P_3^x = P_3$ and therefore $\beta^x = \alpha$.

In conclusion, every element in $H_1 \cap H_Y$ fixes $\{\alpha, \beta\}$ setwise and this completes the proof of the lemma in case (i).

Finally, let us assume (ii) holds, so $a_1 = n/2$, $a_2 \geq 3$ and α and β lie in different parts of X_1 . If $a_2 = n/2$ then the desired conclusion follows from Lemma 3.13, so we may assume that $a_2 < n/2$. As before, set $m = n/a_2$ and note that $m \geq 3$.

Write $P_1 = \{p_1, \dots, p_{n/2}\}$ and $P_2 = \{p_{n/2+1}, \dots, p_n\}$ for the parts of X_1 and assign the colours c_1 and c_2 , respectively. We may assume that $\alpha = p_1$ and $\beta = p_{n/2+1}$. We now

construct an a_2 -partition Y of $[n]$ with parts Q_1, \dots, Q_m and stabiliser H_Y in G , where

$$Q_1 = \{p_1, p_{n/2+2}, \dots, p_{n/2+a_2}\}, \quad Q_2 = \{p_{n/2+1}, p_2, \dots, p_{a_2}\}$$

and we define Q_3, \dots, Q_m by partitioning the remaining points in $[n]$ into subsets of size a_2 with respect to the ordering $p_{a_2+1}, \dots, p_{n/2}, p_{n/2+a_2+1}, \dots, p_n$. To complete the proof, it suffices to show that every element in $H_1 \cap H_Y$ fixes $\{\alpha, \beta\}$ setwise.

To see this, we first claim that Q_2 is the only part of Y which is colour-equivalent to Q_1 (other than Q_1 itself). Seeking a contradiction, suppose that Q_i is colour-equivalent to Q_1 for some $i \geq 3$. From the construction of Y , this is only possible if $n/2 \equiv 1 \pmod{a_2}$, which means that a_2 divides $n - 2$ and n . But this is absurd since $a_2 \geq 3$. This justifies the claim. Similarly, Q_1 is the only part of Y , other than Q_2 , which is colour-equivalent to Q_2 .

Let $x \in H_1 \cap H_Y$ and suppose that $Q_1^x \neq Q_1$. From the previous claim we see that $Q_1^x = Q_2$ and $Q_2^x = Q_1$, which in turn implies that $\alpha^x = \beta$ and $\beta^x = \alpha$ since $a_2 \geq 3$. Finally, let us assume $Q_1^x = Q_1$. Then $Q_2^x = Q_2$ and we deduce that $\alpha^x = \alpha$ and $\beta^x = \beta$. So in all cases, we conclude that x fixes $\{\alpha, \beta\}$ setwise and the proof of the lemma is complete. \square

Remark 3.16. Notice that in Lemma 3.15(ii) we assume the points α and β are contained in different parts of X_1 . Indeed, the conclusion is sometimes false if we drop this condition (for instance, it is straightforward to check that it fails to hold when $a_1 = a_2 = n/2$).

We need one more ingredient before we are in position to prove our main result on imprimitive tuples. This is a special case of a more general result on base sizes for the action of S_n and A_n on partitions (see [12, Proposition 2.4 and Remark 2.8]).

Proposition 3.17. *Let $G = S_n$ or A_n , where $n \geq 8$ is even, and let $H = S_2 \wr S_{n/2}$ be the stabiliser in S_n of a 2-partition of $[n]$. Then there exist $x, y \in G$ such that $H \cap H^x \cap H^y = 1$.*

We are now ready to prove our main result on imprimitive tuples.

Proposition 3.18. *Let $n \geq 6$ be a composite integer. Then*

$$R_{\text{imprim}}(S_n) \leq n - 1, \quad R_{\text{imprim}}(A_n) \leq n - 2.$$

Proof. Let $G = S_n$ or A_n and let $\tau = (J_1, \dots, J_k)$ be a tuple of imprimitive subgroups of G with $k = n - |S_n : G|$. We need to show that τ is regular.

To do this, we may first embed each J_i in a maximal imprimitive subgroup $L_i = S_{a_i} \wr S_{b_i}$ of S_n , where $n = a_i b_i$ and $a_i \geq 2$ for all i , and we may assume that

$$n/2 \geq a_1 \geq a_2 \geq \dots \geq a_{n-2} \geq 2.$$

It suffices to show that there exist elements $g_i \in G$ such that

$$\left(\bigcap_{i=1}^k L_i^{g_i} \right) \cap G = 1. \quad (5)$$

The result for $n \leq 7$ can be checked with the aid of MAGMA, so we will assume $n \geq 8$. Our goal is to prove the following claim:

Claim (\star) . *There exist elements $g_i \in G$ such that $\bigcap_{i=1}^{n-2} L_i^{g_i} \leq \langle (n-1, n) \rangle$.*

Indeed, suppose the claim is true. For $G = A_n$ we have $k = n - 2$, whence (5) holds and the result follows. Now assume $G = S_n$ and let X_{n-1} be any a_{n-1} -partition of $[n]$ with the property that $n - 1$ and n are contained in different parts. Then the stabiliser of X_{n-1} in G is of the form $L_{n-1}^{g_{n-1}}$ for some $g_{n-1} \in G$ and we deduce that (5) holds, which completes the proof of the proposition.

So it just remains to establish Claim (\star) . In view of the transitivity of G on the set of ℓ -partitions of $[n]$ for any ℓ , it suffices to construct a_i -partitions X_i of $[n]$ for $1 \leq i \leq n - 2$ such that

$$H = \bigcap_{i=1}^{n-2} H_i \leq \langle (n-1, n) \rangle,$$

where H_i is the stabiliser of X_i in S_n . (In the course of the proof, we will define various a_i -partitions X_i and we will always write H_i for the stabiliser of X_i in S_n .)

First observe that if $a_{n-4} = 2$, then each L_i with $i \in \{n-4, n-3, n-2\}$ is of the form $S_2 \wr S_{n/2}$, in which case Proposition 3.17 implies that

$$L_{n-4} \cap L_{n-3}^x \cap L_{n-2}^y = 1$$

for some $x, y \in G$. So for the remainder of the proof, we may assume that $a_{n-4} \geq 3$. We now divide the argument into four cases.

Case 1. $a_1 < n/2$

The argument in this case follows very easily from Lemma 3.15. Consider the a_1 -partition

$$X_1 = \{\{1, 2, \dots, a_1\}, \{a_1 + 1, \dots, 2a_1\}, \dots, \{n - a_1 + 1, \dots, n\}\}.$$

Suppose $a_{n-3} \geq 3$. By applying Lemma 3.15(i), we can define an a_2 -partition X_2 such that $H_1 \cap H_2$ fixes $\{1, 2\}$. The same lemma also implies that there is an a_3 -partition X_3 such that $H_2 \cap H_3$ fixes $\{2, 3\}$. And by iteratively applying the same lemma, we can define an a_i -partition X_i such that $H_{i-1} \cap H_i$ fixes $\{i-1, i\}$ for all $4 \leq i \leq n-2$. It follows that $H = \bigcap_{i=1}^{n-2} H_i$ fixes $\{1, \dots, n-2\}$ pointwise and thus $H \leq \langle (n-1, n) \rangle$ as required.

The case $a_{n-3} = 2$ is very similar. By arguing as above, using $a_{n-4} \geq 3$, we can define an a_i -partition X_i for $2 \leq i \leq n-3$ such that $\bigcap_{i=1}^{n-3} H_i$ acts trivially on $\{1, \dots, n-3\}$. We now define the 2-partition

$$X_{n-2} = \{\{1, n-2\}, \{2, n-1\}, \{3, n\}, \{5, 6\}, \dots, \{n-4, n-3\}\}.$$

Since $\bigcap_{i=1}^{n-3} H_i$ fixes 1, 2 and 3, it follows that $\bigcap_{i=1}^{n-2} H_i$ must also fix $n-2$, $n-1$ and n , which forces $\bigcap_{i=1}^{n-2} H_i = 1$.

Case 2. $a_1 = n/2$ and $a_2 < n/2$

Set

$$X_1 = \{\{1, 3, \dots, n/2 + 1\}, \{2, n/2 + 2, \dots, n\}\}$$

and assume $a_{n-3} \geq 3$ for now. Since the points 1 and 2 are contained in different parts of X_1 , Lemma 3.15(ii) implies that we can find an a_2 -partition X_2 such that $H_1 \cap H_2$ fixes $\{1, 2\}$. We can then appeal to Lemma 3.15(i), as in Case 1, to iteratively construct an a_i -partition X_i for $3 \leq i \leq n-2$ such that $\bigcap_{i=2}^{n-2} H_i$ fixes each of the subsets

$$\{2, 3\}, \{3, 4\}, \dots, \{n-3, n-2\}.$$

Consequently, $\bigcap_{i=1}^{n-2} H_i$ fixes $\{1, \dots, n-2\}$ pointwise and the desired result follows.

The case $a_{n-3} = 2$ is very similar. Here we define X_2, \dots, X_{n-3} as above, so that $\bigcap_{i=1}^{n-3} H_i$ fixes $\{1, \dots, n-3\}$ pointwise. By setting

$$X_{n-2} = \{\{1, n-2\}, \{2, n-1\}, \{3, n\}, \{5, 6\}, \dots, \{n-4, n-3\}\}$$

as in Case 1, we deduce that $\bigcap_{i=1}^{n-2} H_i = 1$.

Case 3. $a_1 = a_2 = n/2$ and $a_3 < n/2$

First note that $a_3 \geq 3$ since $n \geq 8$ and $a_{n-4} \geq 3$. We define two $(n/2)$ -partitions

$$\begin{aligned} X_1 &= \{\{1, 4, \dots, n/2 + 2\}, \{2, 3, n/2 + 3, \dots, n\}\} \\ X_2 &= \{\{2, 4, \dots, n/2 + 2\}, \{1, 3, n/2 + 3, \dots, n\}\} \end{aligned}$$

and we note that $H_1 \cap H_2$ fixes $\{1, 2\}$ by Lemma 3.13. In addition, since 2 and 3 are contained in different parts of X_2 , Lemma 3.15(ii) allows us to construct an a_3 -partition X_3 such that $H_2 \cap H_3$ fixes $\{2, 3\}$.

Suppose $a_{n-3} \geq 3$. By applying Lemma 3.15(i), we can iteratively construct an a_i -partition X_i for $4 \leq i \leq n-2$ such that $H_{i-1} \cap H_i$ fixes $\{i-1, i\}$ for all such i . It follows that $\bigcap_{i=1}^{n-2} H_i$ fixes the 2-sets

$$\{1, 2\}, \{2, 3\}, \dots, \{n-3, n-2\}$$

and thus $\bigcap_{i=1}^{n-2} H_i \leq \langle (n-1, n) \rangle$.

Now assume $a_{n-3} = 2$. As above, we can appeal to Lemma 3.15(i) in order to construct partitions X_4, \dots, X_{n-3} such that $H_{i-1} \cap H_i$ fixes $\{i-1, i\}$ for all $i \in \{4, \dots, n-3\}$. Then $\bigcap_{i=1}^{n-3} H_i$ fixes $\{1, \dots, n-3\}$ pointwise and we complete the argument by defining the 2-partition

$$X_{n-2} = \{\{1, n-2\}, \{2, n-1\}, \{3, n\}, \{5, 6\}, \dots, \{n-4, n-3\}\},$$

which forces $\bigcap_{i=1}^{n-2} H_i = 1$.

Case 4. $a_i = n/2$ for some $i \geq 3$

Let $k \in \{1, \dots, n-2\}$ be maximal such that $a_k = n/2$. By appealing to Lemma 3.14, there exist $(n/2)$ -partitions X_1, \dots, X_k such that $\bigcap_{i=1}^k H_i$ fixes $\{1, \dots, k\}$ pointwise. If $k = n-2$ we are done, so we may assume $k \leq n-3$.

Suppose $a_{k+1} = 2$. Here $k+1 \in \{n-3, n-2\}$ and we define

$$X_{k+1} = \{\{1, n-3\}, \{2, n-2\}, \{3, n-1\}, \{4, n\}, \dots, \{n-5, n-4\}\}.$$

Since $n \geq 8$, we note that $\bigcap_{i=1}^k H_i$ fixes 1, 2, 3 and 4, whence $\bigcap_{i=1}^{k+1} H_i = 1$.

Now assume $a_{k+1} \geq 3$. Since k is fixed by $\bigcap_{i=1}^k H_i$, there must exist some $j \in \{1, \dots, k\}$ such that k and $k+1$ lie in different parts of X_j (if not, then $\bigcap_{i=1}^k H_i$ would contain the transposition $(k, k+1)$). Then Lemma 3.15(ii) implies that we can construct an a_{k+1} -partition X_{k+1} such that $H_j \cap H_{k+1}$ fixes $\{k, k+1\}$. If $k = n-3$, then the argument is complete, so we may assume that $k \leq n-4$ for the remainder of the proof.

First assume $a_{n-3} \geq 3$. Here we use Lemma 3.15(i) to construct appropriate partitions X_{k+2}, \dots, X_{n-2} so that $H_{i-1} \cap H_i$ fixes $\{i-1, i\}$ for all $i \in \{k+2, \dots, n-2\}$. It follows that $\bigcap_{i=1}^{n-2} H_i$ fixes the 2-sets $\{k, k+1\}, \dots, \{n-3, n-2\}$ and thus $\bigcap_{i=1}^{n-2} H_i \leq \langle (n-1, n) \rangle$, as claimed.

Finally, suppose $a_{n-3} = 2$. Since $a_{n-4} \geq 3$, we can construct partitions X_{k+2}, \dots, X_{n-3} via Lemma 3.15(i) so that $H_{i-1} \cap H_i$ fixes $\{i-1, i\}$ for all $i \in \{k+2, \dots, n-3\}$. Then $\bigcap_{i=1}^{n-3} H_i$ fixes $\{1, \dots, n-3\}$ pointwise and so by setting

$$X_{n-2} = \{\{1, n-2\}, \{2, n-1\}, \{3, n\}, \{5, 6\}, \dots, \{n-4, n-3\}\}$$

we conclude that $\bigcap_{i=1}^{n-2} H_i = 1$.

This completes the proof of Claim (\star) and the result follows. \square

3.2.3. Proof of Theorem 1. We are now ready to complete the proof of Theorem 1 by showing that $R(S_n) = n-1$ and $R(A_n) = n-2$ as in part (i). To do this, we will bring together our earlier work in Sections 3.1, 3.2.1 and 3.2.2; the key ingredient is the following theorem.

Theorem 3.19. *Let $G = S_n$ or A_n with $n \geq 13$, and let $\tau = (H_1, \dots, H_{n-2})$ be a core-free tuple of subgroups of G . Then there exist elements $g_i \in G$ and $\alpha, \beta \in [n]$ such that*

$$\bigcap_{i=1}^{n-2} H_i^{g_i} \leq \langle (\alpha, \beta) \rangle.$$

Proof. In view of Lemma 2.9, we may assume that $H_i \in \mathcal{M}(G)$ for all i .

Suppose H_1 and H_2 both act primitively on $\{1, \dots, n\}$. Since $n \geq 13$, Theorem 1(ii) implies that $R_{\text{prim}}(S_n) = 2$ and thus $H_1 \cap H_2^{g_2} = 1$ for some $g_2 \in G$. Therefore, for the remainder, we may assume that there is at most one primitive subgroup in τ . By reordering the subgroups, if necessary, we may also assume that if H_i is intransitive, H_j is imprimitive and H_k is primitive, then $i < j$ and $k = n-2$.

Set $\sigma = (a, b, c)$, where a, b and c are the number of intransitive, imprimitive and primitive subgroups in τ , respectively. Then the possibilities for σ are as follows, where $k \geq 1$:

$$(n-2, 0, 0), (0, n-2, 0), (n-3, 0, 1), (0, n-3, 1), (k, n-k-2, 0), (k, n-k-3, 1).$$

We now consider each possibility in turn.

Case 1. $\sigma = (n-2, 0, 0)$ or $(0, n-2, 0)$

First assume $\sigma = (n-2, 0, 0)$. Here τ is an intransitive tuple and this case was handled in the proof of Proposition 3.10 (see (2)). Similarly, τ is an imprimitive tuple if $\sigma = (0, n-2, 0)$ and we can appeal to Claim (\star) in the proof of Proposition 3.18.

Case 2. $\sigma = (n-3, 0, 1)$ or $(0, n-3, 1)$

Next assume $\sigma = (n-3, 0, 1)$, so H_{n-2} is primitive and we have $H_i = (S_{a_i} \times S_{n-a_i}) \cap G$ for $1 \leq i \leq n-3$, where $1 \leq a_i \leq n/2$ and we may assume that $a_1 \leq a_2 \leq \dots \leq a_{n-3}$. For $i \in \{1, \dots, n-3\}$, we define the specific a_i -sets X_i in (1) (see the proof of Lemma 3.9) and we write L_i for the stabiliser of X_i in G . Then $L_i = H_i^{g_i}$ for some $g_i \in G$ and Lemma 3.9 implies that $\bigcap_{i=1}^{n-3} L_i$ fixes $\{1, \dots, n-4\}$ pointwise. If $n \geq 40$, then the lower bound on $\mu(H_{n-2})$ in Theorem 3.3 implies that $\text{supp}(x) \geq 8$ for every nontrivial element $x \in H_{n-2}$, and with the aid of MAGMA it is easy to check that the same conclusion holds for all n in the range $13 \leq n < 40$. This immediately implies that

$$\left(\bigcap_{i=1}^{n-3} L_i \right) \cap H_{n-2} = 1$$

and the result follows.

A very similar argument applies when $\sigma = (0, n-3, 1)$. Here $H_i = (S_{a_i} \wr S_{b_i}) \cap G$ for $i \in \{1, \dots, n-3\}$, where $n = a_i b_i$ and $a_i, b_i \geq 2$. We may assume that $a_1 \geq a_2 \geq \dots \geq a_{n-3}$. By arguing as in the proof of Proposition 3.18, we can construct an a_i -partition X_i for $i \in \{1, \dots, n-3\}$ with stabiliser L_i in G such that $\bigcap_{i=1}^{n-3} L_i$ fixes $\{1, \dots, n-3\}$ pointwise. We now complete the argument as above, appealing to the fact that $\text{supp}(x) \geq 8$ for all $1 \neq x \in H_{n-2}$.

Case 3. $\sigma = (k, n-k-2, 0)$

Here $k \in \{1, \dots, n-3\}$ and recall that we arrange the component subgroups of τ so that for $i \leq k$ we have $H_i = (S_{a_i} \times S_{n-a_i}) \cap G$ with $a_i \leq n/2$, and $H_i = (S_{b_i} \wr S_{c_i}) \cap G$ for $i > k$, where $n = b_i c_i$ and $b_i \geq 2$. In addition, we may assume that $a_1 \leq a_2 \leq \dots \leq a_k$ and $b_{k+1} \geq b_{k+2} \geq \dots \geq b_{n-2}$.

We start by defining the sets X_1, \dots, X_k as in (1) and we write L_i for the stabiliser of X_i in G . Then for each $i \in \{k+1, \dots, n-2\}$ we need to construct a b_i -partition Y_i with stabiliser L_i such that $\bigcap_{i=1}^{n-2} L_i \leq \langle (\alpha, \beta) \rangle$ for some $\alpha, \beta \in [n]$. To do this, we will handle the special case $k \in \{n-4, n-3\}$ separately.

Case 3(a). $k \in \{n-4, n-3\}$

First assume $k = n-3$ and recall that $m = \max\{\alpha : \alpha \in X_k\}$, so $m \in \{n-3, n-2, n-1\}$. Set $L = \bigcap_{i=1}^{n-3} L_i$. Then by applying Lemma 3.9 we deduce that either $L \leq \text{Sym}(X)$ with $X = \{n-3, n-2, n-1\}$ or $\{n-2, n-1, n\}$, or $L \leq \text{Sym}(X) \times \text{Sym}(Y)$ with $X = \{n-3, n-2\}$ and $Y = \{n-1, n\}$.

Suppose $L \leq \text{Sym}(X)$ with $X = \{n-3, n-2, n-1\}$ and let X_{n-2} be a b_{n-2} -partition of $[n]$ such that the part P containing $n-1$ does not contain $n-2$ nor $n-3$. Let L_{n-2} be the stabiliser of X_{n-2} in G . Then L fixes every point in P , with the possible exception of $n-1$, so $\bigcap_{i=1}^{n-2} L_i$ fixes $\{1, \dots, n-4, n-1, n\}$ pointwise and so it is contained in $\langle (n-3, n-2) \rangle$. An entirely similar argument applies if $L \leq \text{Sym}(X)$ with $X = \{n-2, n-1, n\}$, so let us assume $L \leq \text{Sym}(X) \times \text{Sym}(Y)$ with $X = \{n-3, n-2\}$ and $Y = \{n-1, n\}$. Here we choose a b_{n-2} -partition X_{n-2} so that the part containing n does not intersect $\{n-3, n-2, n-1\}$.

If L_{n-2} denotes the stabiliser of X_{n-2} , then by arguing as above, we deduce that $\bigcap_{i=1}^{n-2} L_i$ fixes n , so it must also fix $n-1$ and it is therefore contained in $\langle(n-3, n-2)\rangle$.

A very similar argument applies when $k = n-4$. Set $L = \bigcap_{i=1}^{n-4} L_i$, so Lemma 3.9 implies that either $L \leq \text{Sym}(X)$ with $X = \{n-4, n-3, n-2, n-1\}$ or $\{n-3, n-2, n-1, n\}$, or $L \leq \text{Sym}(X) \times \text{Sym}(Y)$ with $X = \{n-4, n-3, n-2\}$ and $Y = \{n-1, n\}$, or $X = \{n-4, n-3\}$ and $Y = \{n-2, n-1, n\}$.

Suppose $L \leq \text{Sym}(X)$ with $X = \{n-4, n-3, n-2, n-1\}$. Choose a b_{n-3} -partition X_{n-3} with stabiliser L_{n-3} such that the part containing $n-1$ does not intersect $\{n-4, n-3, n-2\}$. Then by arguing as above, we deduce that $\bigcap_{i=1}^{n-3} L_i$ fixes $n-1$ and is therefore contained in $\text{Sym}(Y)$ for $Y = \{n-4, n-3, n-2\}$. We can now complete the argument as in the case $k = n-3$. Similarly, if $L \leq \text{Sym}(X) \times \text{Sym}(Y)$ with $X = \{n-4, n-3, n-2\}$ and $Y = \{n-1, n\}$, then we define a b_{n-3} -partition X_{n-3} so that the part containing n does not meet $\{n-4, n-3, n-2, n-1\}$. Then $\bigcap_{i=1}^{n-3} L_i$ fixes n and is therefore contained in $\text{Sym}(Z)$ for $Z = \{n-4, n-3, n-2\}$. By choosing a b_{n-2} -partition X_{n-2} so that the part containing $n-2$ does not contain $n-4$ nor $n-3$, we conclude that $\bigcap_{i=1}^{n-2} L_i \leq \langle(n-4, n-3)\rangle$, as required.

The other two possibilities (with $k = n-4$) are entirely similar and we omit the details.

Case 3(b). $k \leq n-5$

Now assume $k \leq n-5$. Set $L = \bigcap_{i=1}^k L_i$ and note that L fixes $\{1, \dots, k-1\}$ pointwise by Lemma 3.9. In addition, the same lemma implies that k and n are contained in distinct L -orbits. Therefore, it suffices to construct appropriate partitions X_{k+1}, \dots, X_{n-2} of $[n]$ with stabilisers L_{k+1}, \dots, L_{n-2} such that $\bigcap_{i=k+1}^{n-2} L_i$ fixes $\{k+1, \dots, n-2\}$ pointwise. Indeed, this will imply that $\bigcap_{i=1}^{n-2} L_i \leq \langle g \rangle$, where $g = (k, n-1)$ or $(n-1, n)$.

Suppose $b_{n-4} = 2$, in which case $H_i = (S_2 \wr S_{n/2}) \cap G$ for all $i \in \{n-4, n-3, n-2\}$. Here Proposition 3.17 implies that there exists $g_i \in G$ such that $\bigcap_{i=n-4}^{n-2} H_i^{g_i} = 1$. Therefore, for the remainder of Case 3(b) we may assume that $b_{n-4} \geq 3$.

Let us first consider the special case where $b_i \in \{2, n/2\}$ for all i , in which case $n \geq 14$ is even. Here we have $b_{k+1} = b_{n-4} = n/2$ and we let $j \in \{n-4, n-3, n-2\}$ be maximal such that $b_j = n/2$. We now consider several separate cases.

Suppose $j \leq k+2$. Since $j \geq n-4$, this means that $k \in \{n-6, n-5\}$ and thus $b_{n-2} = 2$. Since $n/2 \geq 7$, we can define a 2-partition X_{n-2} with the property that all of the points k, \dots, n lie in different parts. For $\ell \in \{k, \dots, n\}$, write P_ℓ for the part of X_{n-2} containing ℓ . Since P_ℓ contains at least one point fixed by L , it follows that $L \cap L_{n-2}$ fixes P_ℓ , where L_{n-2} is the stabiliser of X_{n-2} . In turn, this implies that $L \cap L_{n-2}$ fixes every $\ell \in \{k, \dots, n\}$ and thus $L \cap L_{n-2} = 1$.

Now assume $j \geq k+3$. Here Lemma 3.14 implies that we can construct a b_i -partition X_i for $i \in \{k+1, \dots, j\}$ such that $\bigcap_{i=k+1}^j L_i$ fixes $\{k+1, \dots, j\}$ pointwise. If $j = n-2$, then we are done, so let us assume $j = n-4$ or $n-3$. Note that $b_{n-2} = 2$, so $n \geq 14$ is even.

Since $j+1 \geq n-3$, we deduce that $\bigcap_{i=1}^j L_i \leq \text{Sym}(\{k, n-3, \dots, n\})$. Since $n/2 \geq 7$, we can construct a 2-partition X_{j+1} such that each point $\ell \in \{k, n-3, \dots, n\}$ is contained in a distinct part P_ℓ . If L_{j+1} denotes the stabiliser of X_{j+1} , then it is easy to see that $\bigcap_{i=1}^{j+1} L_i = 1$ and the result follows.

To complete the argument in Case 3(b), we may assume $b_r \notin \{2, n/2\}$ for some r . If $b_{n-2} \geq 3$, then we can proceed as in the proof of Proposition 3.18 to produce a b_i -partition X_i for each $i \in \{k+1, \dots, n-2\}$ such that $\bigcap_{i=k+1}^{n-2} L_i$ fixes $\{k+1, \dots, n-2\}$ pointwise, where L_i is the stabiliser of X_i . Finally, suppose $b_{n-2} = 2$ and let $j \in \{n-3, n-2\}$ be minimal such that $b_j = 2$. As in the proof of Proposition 3.18, we can construct a b_i -partition X_i for all $i \in \{k+1, \dots, j\}$ such that $\bigcap_{i=k+1}^j L_i$ fixes $\{k+1, \dots, j\}$ pointwise, whence the desired result follows if $j = n-2$. On the other hand, if $j = n-3$ then $\bigcap_{i=1}^j L_i \leq \text{Sym}(\{k, n-2, n-1, n\})$

and we obtain $\bigcap_{i=1}^{n-2} L_i = 1$ by taking L_{n-2} to be the stabiliser of a 2-partition of $[n]$ such that the points $k, n-2, n-1$ and n all lie in distinct parts.

Case 4. $\sigma = (k, n-k-3, 1)$

In order to complete the proof of the theorem, we may assume $\sigma = (k, n-k-3, 1)$ for some $k \in \{1, \dots, n-4\}$. As before, we order the subgroups in τ so that $H_i = (S_{a_i} \times S_{n-a_i}) \cap G$ for $i \leq k$ and $H_i = (S_{b_i} \wr S_{c_i}) \cap G$ for $i \in \{k+1, \dots, n-3\}$, where $a_i \leq n/2$, $n = b_i c_i$ and $b_i \geq 2$. In addition, H_{n-2} is primitive and we assume that $a_1 \leq a_2 \leq \dots \leq a_k$ and $b_{k+1} \geq b_{k+2} \geq \dots \geq b_{n-3}$. As noted in Case 2, we have $\text{supp}(x) \geq 8$ for all $1 \neq x \in H_{n-2}$. We define the sets X_1, \dots, X_k as in the statement of Lemma 3.9, with stabiliser L_i , and we set $L = \bigcap_{i=1}^k L_i$. By the lemma, we note that L fixes $\{1, \dots, k-1\}$ pointwise.

If $k \geq n-6$ then Lemma 3.9 implies that every element in $L = \bigcap_{i=1}^k L_i$ moves at most 7 points in $[n]$, whence $L \cap H_{n-2} = 1$ and the result follows. Now assume $k \leq n-7$. Here we can proceed as in Case 2 to produce a b_i -partition X_i with stabiliser L_i for $i \in \{k+1, \dots, n-3\}$ such that $\bigcap_{i=k+1}^{n-3} L_i$ fixes $\{k+1, \dots, n-3\}$ pointwise. Therefore, every element in $\bigcap_{i=1}^{n-3} L_i$ moves at most 4 points and we conclude that

$$\left(\bigcap_{i=1}^{n-3} L_i \right) \cap H_{n-2} = 1$$

as required. \square

Remark 3.20. One can check that the conclusion to Theorem 3.19 also holds for $G = S_n$ or A_n with $5 \leq n \leq 12$, with the exception of $G = S_6$. Indeed, if $G = S_6$ and $\tau = (H, H, H, H)$, where $H = S_5$ is primitive, then τ is non-regular and H does not contain a transposition.

Finally, we are now in a position to complete the proof of Theorem 1.

Proof of Theorem 1. It just remains to prove that $R(S_n) \leq n-1$ and $R(A_n) \leq n-2$. The groups with $n \leq 12$ can be handled using MAGMA (see [1] for the details), so we may assume $n \geq 13$. For $G = A_n$, the result now follows immediately from Theorem 3.19, so for the remainder we will assume $G = S_n$. Let $\tau = (H_1, \dots, H_{n-1})$ be a core-free tuple of subgroups of G . By Theorem 3.19, there exists $g_i \in G$ such that $\bigcap_{i=1}^{n-2} H_i^{g_i} \leq \langle (\alpha, \beta) \rangle$ for some $\alpha, \beta \in [n]$, so we just need to show that

$$H_{n-1}^{g_{n-1}} \cap \langle (\alpha, \beta) \rangle = 1$$

for some $g_{n-1} \in G$. This is clear if H_{n-1} is primitive (no core-free primitive subgroup of G contains a transposition). And if $H_{n-1} \leq S_k \times S_{n-k}$ is intransitive, then the result follows because we can always find a k -element subset of $[n]$ that contains α , but not β . Similarly, if $H_{n-1} \leq S_a \wr S_b$ is imprimitive with $a, b \geq 2$, then the result follows because we can find an a -partition of $[n]$ with the property that α and β are contained in distinct parts. This completes the proof of the theorem. \square

With the proof of Theorem 1 in hand, we conclude this section by briefly considering the problem of determining all the non-regular $(n-2)$ -tuples for S_n , and the analogous problem for A_n with respect to $(n-3)$ -tuples. Clearly, if $G = S_n$ and $H = S_{n-1}$ is the stabiliser of a point in $\{1, \dots, n\}$, then the maximal intransitive $(n-2)$ -tuple (H, \dots, H) is non-regular. In the following result, we determine all the maximal intransitive $(n-2)$ -tuples for S_n .

Proposition 3.21. *Let $G = S_n$ with $n \geq 5$ and let τ be a maximal intransitive $(n-2)$ -tuple. Then up to conjugacy and reordering, τ is non-regular if and only if*

$$\tau = (H, \dots, H, K),$$

where $H = S_{n-1}$ and $K = S_k \times S_{n-k}$ for some $1 \leq k \leq n/2$.

Proof. First assume $\tau = (H_1, \dots, H_{n-2})$ has the given form, so $H_{n-2} = S_k \times S_{n-k}$ and $H_i = S_{n-1}$ for all $1 \leq i \leq n-3$. Then for any $g_i \in G$ we observe that

$$\bigcap_{i=1}^{n-3} H_i^{g_i} = \text{Sym}(X),$$

where X is a subset of $[n] = \{1, \dots, n\}$ with $|X| \geq 3$. And so if Y is any subset of $[n]$ of size k , then either $|X \cap Y| \geq 2$ or $|X \cap ([n] \setminus Y)| \geq 2$ and thus $\bigcap_{i=1}^{n-2} H_i^{g_i}$ always contains a transposition. In particular, τ is non-regular.

To complete the proof, suppose $\sigma = (L_1, \dots, L_{n-2})$ is a maximal intransitive $(n-2)$ -tuple, where $L_i = S_{a_i} \times S_{n-a_i}$ and

$$1 \leq a_1 \leq a_2 \leq \dots \leq a_{n-2} \leq n/2$$

with $a_{n-3} \geq 2$. We need to show that σ is regular. To do this, define the a_i -sets X_1, \dots, X_{n-4} as in the statement of Lemma 3.9, with respective stabilisers K_i . Set $H = \bigcap_{i=1}^{n-4} K_i$. Then Lemma 3.9 implies that either

- (a) $H \leq \text{Sym}(X)$ with $X = \{n-4, n-3, n-2, n-1\}$ or $\{n-3, n-2, n-1, n\}$;
- (b) $H \leq \text{Sym}(X) \times \text{Sym}(Y)$ with $X = \{n-4, n-3, n-2\}$ and $Y = \{n-1, n\}$, or $X = \{n-4, n-3\}$ and $Y = \{n-2, n-1, n\}$.

We now define the sets

$$X_{n-3} = \{n-3, n-2, 1, 2, \dots, a_{n-3}-2\}, \quad X_{n-2} = \{n-2, n-1, 1, 2, \dots, a_{n-2}-2\}$$

with stabilisers K_{n-3} and K_{n-2} respectively (so if $a_{n-3} = 2$, then $X_{n-3} = \{n-3, n-2\}$). Then $\bigcap_{i=1}^{n-2} K_i = 1$ and we are done. \square

Remark 3.22. For $G = A_n$, an entirely similar argument shows that the maximal intransitive $(n-3)$ -tuples for G are precisely those of the form (H, \dots, H, K) , up to conjugacy and reordering, where $H = A_{n-1}$ and $K = (S_k \times S_{n-k}) \cap G$ for some $1 \leq k \leq n/2$.

Remark 3.23. It would be interesting to extend Proposition 3.21 by giving a complete classification of all the non-regular $(n-2)$ -tuples for $G = S_n$. Notice that if $n = 2m$ is even, $H = S_{n-1}$ and $K = S_m \wr S_2$, then it is easy to see that (H, \dots, H, K) is a non-regular $(n-2)$ -tuple with a transitive component subgroup. And by replacing K by the index-two subgroup $L = S_m \times S_m$, we obtain a non-regular $(n-2)$ -tuple containing a non-maximal subgroup. For $n \geq 5$ odd, we are not aware of any non-regular $(n-2)$ -tuples for G that are not of the form described in Proposition 3.21.

4. SPORADIC GROUPS

For the remainder of the paper, we assume G is an almost simple sporadic group with socle T and our aim is to prove Theorem 2. We begin in Section 4.1 by computing the exact regularity number of G , which establishes part (i) of Theorem 2. We then prove part (ii) in Section 4.2, where we show that every soluble triple of subgroups of G is regular. We apply computational methods throughout this section, working extensively with MAGMA [5] and GAP [28]. We refer the reader to Section 2.4 for a brief overview of some of the main methods, which complements the more detailed discussion in the supplementary file [1].

As explained in Wilson's survey article [50], the maximal subgroups of G are known up to conjugacy, apart from the problem of determining the status of a handful of candidate almost simple maximal subgroups of the Monster \mathbb{M} (and the latter ambiguity appears to have been recently resolved by Dietrich et al. in [25]). In addition, the character table of G is available in the GAP Character Table Library [6], together with the character tables of all the maximal subgroups of G and the associated fusion maps on conjugacy classes (excluding $G = \mathbb{M}$). A great deal of useful information about the sporadic groups is available in the Web Atlas [49].

4.1. The regularity number. Let G be an almost simple sporadic group with socle T , so $G = T$ or $T.2$. Recall that we define $\mathcal{M}(G)$ to be the set of core-free subgroups of G that are maximal in either G or T (so if $G = T$, then $\mathcal{M}(G)$ is just the set of maximal subgroups of G). As noted in Remark 2.10, if $G = T.2$ and $H < T$ is maximal with $N_G(H) \neq H$, then H is contained in a core-free maximal subgroup of G . Therefore, if we define $\mathcal{M}'(G) = \mathcal{A} \cup \mathcal{B}$, where

$$\begin{aligned} \mathcal{A} &= \{\text{core-free maximal subgroups of } G\} \\ \mathcal{B} &= \{\text{maximal subgroups } H \text{ of } T \text{ with } N_G(H) = H\} \end{aligned}$$

then

$$B(G) = \max\{b(G, H) : H \in \mathcal{M}'(G)\}$$

and $R(G)$ is the smallest positive integer k such that every k -tuple of subgroups in $\mathcal{M}'(G)$ is regular. For the remainder of this section, it will be convenient to work with the following definition (note that if $G = T$, then every *large* tuple is *maximal*, and vice versa).

Definition 4.1. A k -tuple of subgroups (H_1, \dots, H_k) of G is *large* if each component subgroup H_i is contained in the set $\mathcal{M}'(G)$ defined above.

Recall that $B(G) \geq 3$ by Proposition 2.7. We begin by recording the exact base number, which is easily obtained from [16].

Proposition 4.2. *Let G be an almost simple sporadic group with socle T . Then the base number $B(G)$ is recorded in Table 3.*

Proof. If $G = T$, then we can immediately read off $B(G)$ from the main theorem of [16], so let us assume $G = T.2$ and $H \in \mathcal{M}'(G)$. If H is maximal in G , then $b(G, H)$ is computed in [16], so we may assume that $H < T$ is maximal and $N_G(H) = H$. Here $b(T, H)$ is recorded in [16], which is clearly an upper bound for $b(G, H)$.

By inspecting [16], we see that $G = M_{12}.2$ is the only group that arises where $B(T)$ is greater than the maximal value of $b(G, H)$ over all core-free maximal subgroups H of G . Here it is easy to check that if $H = M_{11} < T$, then $b(G, H) = 5$ and we deduce that $B(G) = 5$. \square

Now let us turn to the regularity number of G . Our main result is the following, which establishes part (i) of Theorem 2. See Remark 4.5 below for comments on the information recorded in the final column of Table 3. In (ii), we write \mathbb{B} and \mathbb{M} for the Baby Monster and Monster groups, respectively.

Proposition 4.3. *Let G be an almost simple sporadic group with socle T .*

- (i) *The regularity number $R(G) = k$ is recorded in Table 3.*
- (ii) *For $G \neq \mathbb{B}, \mathbb{M}$, a complete list of the large non-regular $(k - 1)$ -tuples, up to ordering and conjugacy, is presented in the final column of Table 3.*

As an immediate corollary, we obtain a strong form of Conjecture 1(ii) for almost simple sporadic groups.

Corollary 4.4. *We have $R(G) \leq 7$, with equality if and only if $G = M_{24}$. Moreover, if $G = M_{24}$ then a 6-tuple is non-regular if and only if every component subgroup is conjugate to M_{23} .*

Remark 4.5. Suppose $G \neq \mathbb{B}, \mathbb{M}$ and $R(G) = k$. In the final column of Table 3 we list all the large non-regular $(k - 1)$ -tuples for G , up to conjugacy and ordering. To do this, we use the notation $(a_1, \dots, a_{k-1}) \in \mathbb{N}^{k-1}$ to encode each tuple $\tau = (H_1, \dots, H_{k-1})$, noting that the corresponding subgroups are recorded in Tables 4 and 5. So for example, if $G = J_3.2$, then $(H_1, H_2) = (L_2(17) \times 2, (3 \times M_{10}):2)$ is the non-regular pair labelled (3, 4) in Table 3. We refer the reader to Remark 4.11 for comments on the groups \mathbb{B} and \mathbb{M} we are excluding here.

G	$B(G)$	$R(G)$
M_{11}	4	5 (1, 1, 2, 2), (1, 2, 2, 2)
M_{12}	5	6 (1, 1, 1, 2, 2), (1, 1, 2, 2, 2)
$M_{12.2}$	5	5 (1, 1, 1, 1), (1, 1, 1, 2)
M_{22}	5	5 (1, 1, 1, 1), (1, 1, 1, 2)
$M_{22.2}$	5	5 (1, 1, 1, i), (1, 1, 2, 2), (1, 1, 2, 4), $1 \leq i \leq 5$
M_{23}	6	6 (1, 1, 1, 1, 1)
M_{24}	7	7 (1, 1, 1, 1, 1, 1)
J_1	3	3 (1, 1), (1, 2)
J_2	4	4 (1, 1, 1)
$J_{2.2}$	4	4 (1, 1, i), $1 \leq i \leq 4$
J_3	3	3 (1, i), $1 \leq i \leq 8$
$J_{3.2}$	3	3 (1, i), (3, 4), (3, 6), (4, 4), $1 \leq i \leq 7$
HS	4	4 (1, 1, 1, i), (1, 1, 2, 2), (1, 1, 3, 3), $1 \leq i \leq 3$
HS.2	5	5 (1, 1, 1, i), $1 \leq i \leq 4$
McL	5	5 (1, 1, 1, 1)
McL.2	5	5 (1, 1, 1, 1)
He	4	4 (1, 1, 1), (1, 1, 2)
He.2	4	4 (1, 1, 1), (1, 1, 2)
Suz	4	4 (1, 1, i), (1, 2, 2), (1, 2, 3), (1, 3, 3), $1 \leq i \leq 7$
Suz.2	4	4 (1, 1, i), (1, 2, 2), (1, 2, 3), (1, 3, 3), $1 \leq i \leq 9$
O'N	3	3 (1, i_1), (2, i_2), $j \leq i_j \leq 4$
O'N.2	3	3 (1, i), (2, 3), (3, 3), $1 \leq i \leq 3$
HN	3	3 (1, i_1), (2, i_2), (3, 3), $j \leq i_j \leq 12$
HN.2	3	3 (1, i), (2, j), (2, 12), (3, 3), (3, 4), $1 \leq i \leq 12$, $2 \leq j \leq 10$
Co ₃	6	6 (1, 1, 1, 1, 1)
Co ₂	6	6 (1, 1, 1, 1, 1)
Co ₁	5	5 (1, 1, 1, 1)
Ru	4	4 (1, 1, 1)
Fi ₂₂	5	5 (1, 1, 1, 1), (1, 1, 1, 2)
Fi _{22.2}	6	6 (1, 1, 1, 1, 1)
Fi ₂₃	5	5 (1, 1, 1, 1), (1, 1, 1, 2), (1, 1, 2, 2), (1, 2, 2, 2)
Fi' ₂₄	5	5 (1, 1, 1, 1), (1, 1, 1, 2)
Fi ₂₄	5	5 (1, 1, 1, i), $1 \leq i \leq 3$
J ₄	3	3 (1, i), (2, j), (3, 3), $1 \leq i \leq 5$, $2 \leq j \leq 4$
Ly	3	3 (1, i_1), (2, i_2), $j \leq i_j \leq 6$
Th	3	3 (1, i), (2, 2), $1 \leq i \leq 4$
\mathbb{B}	4	4 See Remark 4.11
\mathbb{M}	3	3 See Remark 4.11

TABLE 3. The base and regularity numbers for sporadic groups

Let G be an almost simple sporadic group with socle T . Our proof of Proposition 4.3 is entirely computational, working with MAGMA (version 2.28-4) [5] and GAP (version 4.11.1) [28]. In particular, the information available in the GAP Character Table Library [6] is essential for handling some of the larger groups. We begin by dividing the 26 possibilities for T into the following four collections, and we will handle each one in turn:

$$\mathcal{A}_1 = \{M_{11}, M_{12}, M_{22}, M_{23}, M_{24}, J_1, J_2, J_3, HS, McL, He, Suz, O'N, Ru, Co_3, Co_2, Fi_{22}, Fi_{23}\}$$

$$\mathcal{A}_2 = \{Co_1, HN, Fi'_{24}\}$$

$$\mathcal{A}_3 = \{J_4, Ly, Th\}$$

$$\mathcal{A}_4 = \{\mathbb{B}, \mathbb{M}\}$$

Lemma 4.6. *The conclusion to Proposition 4.3 holds for the groups with $T \in \mathcal{A}_1$.*

Proof. We begin by reading off $B(G) = b$ from Proposition 4.2 and we fix a large $(b-1)$ -tuple $\sigma = (H_1, \dots, H_{b-1})$. As noted in Lemma 2.6, if

$$\prod_{i=1}^{b-1} |H_i| > |G|^{b-2},$$

	1	2	3	4	5	6
M ₁₁	M ₁₀	L ₂ (11)				
M ₁₂	M ₁₁	M ₁₁				
M ₁₂ .2	M ₁₁	A ₆ .2 ²				
M ₂₂	L ₃ (4)	2 ⁴ :A ₆				
M ₂₂ .2	L ₃ (4).2	2 ⁴ :S ₆	2 ⁵ :S ₅	2 ³ :L ₃ (2) × 2	A ₆ .2 ²	
M ₂₃	M ₂₂					
M ₂₄	M ₂₃					
J ₁	L ₂ (11)	19:6				
J ₂	U ₃ (3)					
J ₂ .2	U ₃ (3):2	3.A ₆ .2 ²	2 ¹⁺⁴ .S ₅	2 ²⁺⁴ :(3 × S ₃).2		
J ₃	L ₂ (16):2	L ₂ (19)	L ₂ (19)	2 ⁴ :(3 × A ₅)	L ₂ (17)	(3 × A ₆).2
J ₃ .2	L ₂ (16):4	2 ⁴ :(3 × A ₅).2	L ₂ (17) × 2	(3 × M ₁₀):2	3 ²⁺¹⁺² :8.2	2 ¹⁺⁴ :S ₅
HS	M ₂₂	U ₃ (5):2	U ₃ (5):2			
HS.2	M ₂₂ .2	L ₃ (4):2 ²	S ₈ × 2	U ₃ (5):2		
McL	U ₄ (3)					
McL.2	U ₄ (3):2					
He	Sp ₄ (4):2	2 ² .L ₃ (4).S ₃				
He.2	Sp ₄ (4):4	2 ² .L ₃ (4).D ₁₂				
Suz	G ₂ (4)	3.U ₄ (3).2	U ₅ (2)	2 ¹⁺⁶ .U ₄ (2)	3 ⁵ :M ₁₁	J ₂ :2
Suz.2	G ₂ (4):2	3.U ₄ (3).2 ²	U ₅ (2):2	2 ¹⁺⁶ .U ₄ (2).2	3 ⁵ :(M ₁₁ × 2)	J ₂ :2 × 2
O'N	L ₃ (7):2	L ₃ (7):2	J ₁	4.L ₃ (4):2		
O'N.2	L ₃ (7):2	7 ¹⁺² :(3 × D ₁₆)	J ₁ × 2			
HN	A ₁₂	2.HS.2	U ₃ (8):3	2 ¹⁺⁸ .(A ₅ × A ₅).2	(D ₁₀ × U ₃ (5)).2	5 ¹⁺⁴ .2 ¹⁺⁴ .5.4
HN.2	S ₁₂	4.HS.2	U ₃ (8):6	2 ¹⁺⁸ .(A ₅ × A ₅).2.2	5:4 × U ₃ (5)	5 ¹⁺⁴ .2 ¹⁺⁴ .5.4.2
Co ₃	McL.2					
Co ₂	U ₆ (2):2					
Co ₁	Co ₂					
Ru	² F ₄ (2)					
Fi ₂₂	2.U ₆ (2)	2 ¹⁰ .M ₂₂				
Fi ₂₂ .2	2.U ₆ (2).2					
Fi ₂₃	2.Fi ₂₂	PΩ ₈ ⁺ (3):S ₃				
J ₄	2 ¹¹ :M ₂₄	2 ¹⁺¹² .3.M ₂₂ :2	2 ¹⁰ :L ₅ (2)	2 ³⁺¹² .(S ₅ × L ₃ (2))	U ₃ (11):2	
Ly	G ₂ (5)	3.McL:2	5 ³ .L ₃ (5)	2.A ₁₁	5 ¹⁺⁴ .4.S ₆	3 ⁵ :(2 × M ₁₁)
Th	³ D ₄ (2):3	2 ⁵ .L ₅ (2)	2 ¹⁺⁸ .A ₉	U ₃ (8):6		
Fi ₂₄ '	Fi ₂₃	2.Fi ₂₂ .2				
Fi ₂₄	Fi ₂₃ × 2	(2 × 2.Fi ₂₂):2	S ₃ × PΩ ₈ ⁺ (3):S ₃			

TABLE 4. The subgroups recorded in Table 3, Part I

	7	8	9	10	11	12
M ₁₂ .2	S ₄ × S ₃	S ₅				
J ₃	3 ²⁺¹⁺² :8	2 ¹⁺⁴ :A ₅				
J ₃ .2	L ₂ (19)					
Suz	2 ²⁺⁸ :(A ₅ × S ₃)					
Suz.2	2 ⁴⁺⁶ :3.S ₆	(A ₄ × L ₃ (4):2):2	2 ²⁺⁸ :(S ₅ × S ₃)			
HN	2 ⁶ .U ₄ (2)	(A ₆ × A ₆).D ₈	2 ³⁺²⁺⁶ .(3 × L ₃ (2))	5 ²⁺¹⁺² .4.A ₅	M ₁₂ .2	M ₁₂ .2
HN.2	2 ⁶ .U ₄ (2).2	(S ₆ × S ₆).2 ²	2 ³⁺²⁺⁶ .(3 × L ₃ (2)).2	5 ²⁺¹⁺² .4.A ₅ .2	3 ⁴ :2.(S ₄ × S ₄).2	M ₁₂ .2

TABLE 5. The subgroups recorded in Table 3, Part II

then G does not have a regular orbit on $G/H_1 \times \cdots \times G/H_{b-1}$ and thus σ is non-regular, so let us assume otherwise.

We use the MAGMA functions `AutomorphismGroupSimpleGroup` and `MaximalSubgroups` to construct G and each H_i as subgroups of S_n , where $n = m(G)$ is the minimal degree of a faithful primitive permutation representation of G . We can then randomly search for elements $g_i \in G$ such that $\bigcap_i H_i^{g_i} = 1$. If our search is successful after a specified number of attempts, then σ is regular and we repeat for the next $(b-1)$ -tuple in our list. In every case, this process quickly produces a distinguished collection of large $(b-1)$ -tuples that require closer inspection in order to determine their regularity status (of course, there has to be at least one such tuple since $B(G) = b$).

Let $\sigma = (H_1, \dots, H_{b-1})$ be a large tuple for which random search is inconclusive. Now σ is regular if and only if H_1 has a regular orbit on $Y = G/H_2 \times \dots \times G/H_{b-1}$, and we can use MAGMA to determine whether or not such an orbit exists. To do this, we first reorder the subgroups in σ so that $|H_1|$ is minimal. We then use the function `CoSetAction` to construct G as a permutation group on each set $\Gamma_i = G/H_i$ with $2 \leq i \leq b-1$ and we compute a set of orbit representatives $\{\alpha_1, \dots, \alpha_m\}$ for the action of H_1 on Γ_2 , with corresponding stabilisers $L_i = (H_1)_{\alpha_i}$. If $b = 3$ then σ is regular if and only if at least one of these stabilisers is trivial. And if $b = 4$, then we can repeat the process; for each i , we find a set of orbit representatives for the action of L_i on Γ_3 and we determine if any of the corresponding stabilisers in L_i are trivial. And similarly for $b \geq 5$. In this way, we obtain the complete list of large non-regular $(b-1)$ -tuples, up to ordering and conjugacy.

Next observe that a b -tuple $(H_1, \dots, H_{b-1}, H_b)$ is non-regular only if (H_1, \dots, H_{b-1}) is non-regular, so we can work through our list of non-regular $(b-1)$ -tuples, proceeding as above, to determine whether or not there are any non-regular b -tuples.

For $G \neq M_{11}, M_{12}$, we find that there are no such tuples and we conclude that $R(G) = B(G) = b$. In addition, the list of large non-regular $(b-1)$ -tuples is presented in the final column of Table 3 (see Remark 4.5). However, large non-regular b -tuples do exist when $G = M_{11}$ or M_{12} ; it is easy to determine all such tuples, and it is also easy to check that every $(b+1)$ -tuple is regular. So in these two cases, we deduce that $R(G) = B(G) + 1$. \square

Lemma 4.7. *The conclusion to Proposition 4.3 holds for the groups with $T \in \mathcal{A}_2$.*

Proof. First assume $G = \text{Co}_1$, so $B(G) = 5$. Here it is possible to proceed as in Case 1, but there is a more efficient way to handle the computation, which we can also implement in some of the other remaining cases.

Let $\sigma = (H_1, \dots, H_4)$ be a maximal 4-tuple and define $\widehat{Q}(G, \sigma)$ as in Lemma 2.1. Recall that σ is regular if $\widehat{Q}(G, \sigma) < 1$. The character tables of G and each maximal subgroup H_i are available in the GAP Character Table Library [6], together with the fusion map from H_i -classes to G -classes. This allows us to compute

$$\text{fpr}(x, G/H_i) = \frac{|x^G \cap H_i|}{|x^G|}$$

for every element $x \in G$, which in turn allows us to calculate $\widehat{Q}(G, \sigma)$. In this way, we can determine all the maximal 4-tuples σ with $\widehat{Q}(G, \sigma) \geq 1$, up to conjugacy and ordering. The complete list is as follows:

$$(1, 1, 1, 1), (1, 1, 1, 2), (1, 1, 1, 3), (1, 1, 1, 4), (1, 1, 1, 5), (1, 1, 1, 6),$$

corresponding to the labelling

$$1: \text{Co}_2, \quad 2: 3.\text{Suz}:2, \quad 3: 2^{11}:\text{M}_{24}, \quad 4: \text{Co}_3, \quad 5: 2^{1+8}.\Omega_8^+(2), \quad 6: \text{U}_6(2):S_3$$

Suppose $\sigma = (H_1, \dots, H_4)$ is a maximal 4-tuple with $\widehat{Q}(G, \sigma) \geq 1$, where we have $H_i = \text{Co}_2$ for $1 \leq i \leq 3$. As before, we can use MAGMA to construct G and each component H_i as subgroups of S_n (with $n = |G : H_1| = 98280$) and we can use random search to show that the tuples $(1, 1, 1, j)$ with $2 \leq j \leq 6$ are regular. By [16] we know that $b(G, H_1) = 5$, so $(1, 1, 1, 1)$ is non-regular. Finally, for each representative H of a conjugacy class of maximal subgroups of G , we define $\tau = (H_1, \dots, H_4, H)$ with $H_i = \text{Co}_2$ for all i and then using GAP we find that $\widehat{Q}(G, \tau) < 1$. This allows us to conclude that every maximal 5-tuple is regular and thus $R(G) = B(G) = 5$.

We can handle the groups $G = \text{HN}$ and $\text{HN}.2$ in a similar fashion. In both cases we have $B(G) = 3$ and we can use GAP to determine all the large pairs σ with $\widehat{Q}(G, \sigma) \geq 1$. This yields a short list of candidate non-regular pairs and it is straightforward to show that $\widehat{Q}(G, \tau) < 1$ for every large triple τ , whence $R(G) = B(G) = 3$.

To complete the proof in these two cases, we inspect each large pair $\sigma = (H_1, H_2)$ with $\widehat{Q}(G, \sigma) \geq 1$. As before, if $|H_1||H_2| > |G|$ then σ is non-regular. In order to handle the remaining cases, we first use MAGMA to construct G , H_1 and H_2 as subgroups of S_n with $n = 1140000$. In several cases, we can then use random search to find an element $x \in G$ with $H_1 \cap H_2^x = 1$, which allows us to conclude that σ is regular.

For example, if $G = \text{HN.2}$ then this approach reduces the problem to determining whether or not the following pairs are regular:

$$(1, 11), (1, 12), (1, 13), (2, 10), (2, 11), (2, 12), (2, 13), (3, 4)$$

with respect to the corresponding subgroups:

$$\begin{array}{llll} 1: S_{12}, & 2: 4.\text{HS}.2, & 3: \text{U}_3(8):6, & 4: 2^{1+8}:(A_5 \times A_5).2^2 \\ 10: 5^{2+1+2}.4.A_5.2, & 11: 3^4.2.(S_4 \times S_4).2, & 12: \text{M}_{12}.2, & 13: 3^{1+4}.4.S_5 \end{array}$$

Let K_i be the subgroup numbered i in this list. Since the given permutation representation of G (of degree 1140000) corresponds to the action of G on the cosets of $K_1 = S_{12}$, it is easy to show that K_{13} has a regular orbit on G/K_1 , but K_{11} and K_{12} do not. So $(1, 13)$ is regular and $(1, 11), (1, 12)$ are non-regular. Similarly, using `CosetAction` we can show that K_{11} and K_{12} have regular orbits on G/K_2 , but K_{10} and K_{13} do not. In the same way, we deduce that K_4 does not have a regular orbit on G/K_3 . Therefore, $(2, 11)$ and $(2, 12)$ are regular, whereas $(2, 10), (2, 13)$ and $(3, 4)$ are non-regular. The case $G = \text{HN}$ is very similar.

Next assume $G = \text{Fi}'_{24}$, so $B(G) = 5$ and we claim that $R(G) = 5$. To see this, we first use the GAP Character Table Library to identify all the maximal 4-tuples σ with $\widehat{Q}(G, \sigma) \geq 1$. The tuples that arise are of the form $(1, 1, 1, i)$ with $1 \leq i \leq 6$, where the component subgroups are labelled as follows:

$$1: \text{Fi}_{23}, \quad 2: 2.\text{Fi}_{22}.2, \quad 3: (3 \times \text{P}\Omega_8^+(3).3).2, \quad 4: \Omega_{10}^-(2), \quad 5: 3^7.\Omega_7(3), \quad 6: 3^{1+10}:\text{U}_5(2).2$$

Therefore, any non-regular maximal 5-tuple must be of the form

$$\tau = (H_1, H_2, H_3, H_4, H_5)$$

(up to conjugacy and ordering), where $H_1 = H_2 = H_3 = \text{Fi}_{23}$ and H_4 is one of the six subgroups listed above. Using GAP we find that $\widehat{Q}(G, \tau) \geq 1$ if and only if $H_4 = H_5 = \text{Fi}_{23}$, which is regular since $B(G) = 5$. Therefore $R(G) = B(G) = 5$ and it just remains to determine all the non-regular maximal 4-tuples.

To do this, we return to the above list of maximal 4-tuples σ with $\widehat{Q}(G, \sigma) \geq 1$, noting that $(1, 1, 1, 1)$ is non-regular by [16]. To handle the remaining possibilities, we first use MAGMA to construct G as a permutation group of degree 306936, which corresponds to the action of G on the cosets of Fi_{23} . The command `MaximalSubgroups` is not effective, but we can construct representatives of the relevant conjugacy classes of maximal subgroups using explicit generators in the Web Atlas [49], which are given as words in the standard generators for G (more precisely, we work in $L = G.2 = \text{Fi}_{24}$ and we use explicit generators to construct $N_L(H)$, intersecting with G to get H). Then random search shows that each 4-tuple $(1, 1, 1, i)$ with $3 \leq i \leq 6$ is regular.

Finally, suppose $\sigma = (H, H, H, K)$, where $H = \text{Fi}_{23}$ and $K = 2.\text{Fi}_{22}.2$. Here we can work directly with the permutation representation of G of degree 306936, which corresponds to the action of G on $\Gamma = G/H$. It is straightforward to compute the order of every 3-point stabiliser with respect to the action of K on Γ and we find that every such stabiliser is nontrivial. This immediately implies that σ is non-regular and this completes the proof for $G = \text{Fi}'_{24}$. The case $G = \text{Fi}_{24}$ is entirely similar. \square

Next we turn to the groups in \mathcal{A}_3 . We begin by establishing the non-regularity of a certain collection of maximal pairs.

Lemma 4.8. *All of the maximal pairs (H, K) in Table 6 are non-regular.*

G	H	K	$ G : H $
Ly	$G_2(5)$	$3^5:(2 \times M_{11})$	8835156
	3.McL.2	$5^{1+4}:4S_6$	9606125
	3.McL.2	$3^5:(2 \times M_{11})$	9606125
Th	${}^3D_4(2):3$	$2^{1+8}.A_9$	143127000
	${}^3D_4(2):3$	$U_3(8):6$	143127000
J ₄	$2^{11}.M_{24}$	$U_3(11).2$	173067389
	$2^{1+12}.3.M_{22}.2$	$2^{3+12}.(S_5 \times L_3(2))$	3980549947

TABLE 6. The non-regular pairs in Lemma 4.8

Proof. First observe that $|H||K| < |G|$ in every case, so we cannot appeal to Lemma 2.6. Using MAGMA, we can construct G , H and K as subgroups of the matrix group $GL_d(p)$, where $(d, p) = (111, 5)$, $(248, 2)$ and $(112, 2)$ for $G = \text{Ly}$, Th and J_4 , respectively. This allows us to randomly search for an element $g \in G$ with $H \cap K^g = 1$, but this search is inconclusive in every case. In addition, the size of $|G : H|$ is prohibitive and we are not able to use `CosetAction` to construct G as a permutation group on G/H , as we have done in previous cases.

We thank Derek Holt (personal communication) for resolving the first 6 cases in Table 6. To illustrate his approach, let us assume $G = \text{Ly}$, $H = G_2(5)$ and $K = 3^5:(2 \times M_{11})$, as in the first row of the table, and consider the irreducible module $V = \mathbb{F}_5^{111}$ for G . Holt proceeds by identifying an irreducible 7-dimensional submodule W for H , which he then uses, via the function `OrbitImage`, to construct G as a permutation group on G/H . Then as before, he shows that (H, K) is non-regular by determining all of the K -orbits on G/H . The other 5 cases are handled in a similar fashion.

Finally, suppose $G = \text{J}_4$ and $\tau = (H, K)$, where

$$H = 2^{1+12}.3.M_{22}.2, \quad K = 2^{3+12}.(S_5 \times L_3(2)).$$

This case is more difficult since the index $|G : H| = 3980549947$ is very large. We are grateful to Jürgen Müller (personal communication) for resolving this case (which was later independently verified by Holt). Müller did this by using the GAP package ORB [44] to establish the existence of 54 non-regular K -orbits on $\Gamma = G/H$ (via random search), labelled $\Gamma_1, \dots, \Gamma_{54}$, with the property that

$$|G : H| - \sum_{i=1}^{54} |\Gamma_i| = 3980549947 - 3980074560 < |K|.$$

This immediately implies that K has no regular orbit on G/H , so (H, K) is non-regular and the proof of the lemma is complete. \square

Lemma 4.9. *The conclusion to Proposition 4.3 holds for the groups with $T \in \mathcal{A}_3$.*

Proof. Here $G = \text{Ly}$, Th or J_4 , with $B(G) = 3$ for each group. As in previous cases, we can use GAP to show that $\widehat{Q}(G, \tau) < 1$ for every maximal triple τ , whence $R(G) = B(G) = 3$ and it just remains to determine all the non-regular maximal pairs.

All three cases are very similar, so for brevity we will assume $G = \text{Th}$. As usual, we begin by using GAP to find all the maximal pairs $\sigma = (H_1, H_2)$ with $\widehat{Q}(G, \sigma) \geq 1$, which yields the following list:

$$(1, i), (2, 2), (2, 3), (3, 3)$$

with $1 \leq i \leq 5$. Here the relevant subgroups are as follows:

$$1: {}^3D_4(2):3, \quad 2: 2^5.L_5(2), \quad 3: 2^{1+8}.A_9, \quad 4: U_3(8):6, \quad 5: (3 \times G_2(3)):2.$$

From [16], we see that $(1, 1)$ and $(2, 2)$ are non-regular, whereas $(3, 3)$ is regular. And by computing orders, we deduce that $(1, 2)$ is non-regular. We can then use MAGMA and random search to show that $(1, 5)$ and $(2, 3)$ are regular, working inside the matrix group $GL_{248}(2)$. Finally, we note that the two remaining pairs $(1, 3)$ and $(1, 4)$ were handled in Lemma 4.8. \square

In order to complete the proof of Proposition 4.3, we may assume $G = \mathbb{B}$ or \mathbb{M} .

Lemma 4.10. *The conclusion to Proposition 4.3 holds for the groups with $T \in \mathcal{A}_4$.*

Proof. First assume $G = \mathbb{B}$ is the Baby Monster and note that $B(G) = 4$. As before, the character table of G and every maximal subgroup of G is available in the GAP Character Table Library [6]. Moreover, if H is a maximal subgroup, then the fusion map from H -classes to G -classes is also available unless $H = (2^2 \times F_4(2)):2$. In order to handle the latter case, we can use the function `PossibleClassFusions` to determine a list of 64 candidate fusion maps and we find that each fixed point ratio $\text{fpr}(x, G/H)$ is independent of the choice of map. Therefore, we can compute $\widehat{Q}(G, \tau)$ precisely for every maximal k -tuple τ and we can now proceed as we have done in previous cases.

First we use GAP to determine the list of maximal triples σ with $\widehat{Q}(G, \sigma) \geq 1$:

$$(1, 1, i), (1, 1, 6), (1, 1, 7), (1, 2, 2)$$

where $1 \leq i \leq 4$ and the relevant maximal subgroups are labelled as follows:

$$\begin{array}{llll} 1: 2^2.E_6(2):2, & 2: 2^{1+22}.Co_2, & 3: Fi_{23}, & 4: 2^{9+16}.Sp_8(2), \\ 5: Th, & 6: (2^2 \times F_4(2)):2, & 7: 2^{2+10+20}.(M_{22}:2 \times S_3) \end{array} \quad (6)$$

Working with this list of candidate non-regular triples, it is easy to check that $\widehat{Q}(G, \tau) < 1$ for every maximal 4-tuple τ and thus $R(G) = B(G) = 4$. We refer the reader to Remark 4.11(a) for comments on the problem of determining all the non-regular maximal triples when $G = \mathbb{B}$.

Finally, let us assume $G = \mathbb{M}$ is the Monster, so $B(G) = 3$ by [16]. We claim that $\widehat{Q}(G, \tau) < 1$ for every maximal triple τ , which implies that $R(G) = B(G) = 3$.

First recall that G has 46 conjugacy classes of maximal subgroups (see the main theorem of [25]). For 31 of these classes, the character table of a representative H and the fusion map from H -classes to G -classes is available in the GAP Character Table Library [6] via the function `NamesOfFusionSources`. The relevant subgroups are as follows:

$$\begin{array}{llll} 2.\mathbb{B} & 2^{1+24}.Co_1 & 3.Fi_{24} & 2^2.2E_6(2):S_3 \\ 3^{1+12}.2.Suz:2 & S_3 \times Th & (D_{10} \times HN).2 & 5^{1+6}.2.J_2.4 \\ (7:3 \times He):2 & (A_5 \times A_{12}):2 & 5^{3+3}.(2 \times L_3(5)) & (A_6)^3.(2 \times S_4) \\ (A_5 \times U_3(8):3):2 & 5^{2+2+4}.(S_3 \times GL_2(5)) & (L_3(2) \times Sp_4(4):2).2 & 7^{1+4}.(3 \times 2S_7) \\ (5^2:[2^4] \times U_3(5)).S_3 & (L_2(11) \times M_{12}):2 & (A_7 \times (A_5 \times A_5):2^2):2 & 5^4:(3 \times 2.L_2(25)).2 \\ 7^{2+1+2}.GL_2(7) & M_{11} \times A_6.2^2 & (S_5)^3.S_3 & 13^2.2.L_2(13).4 \\ (7^2:(3 \times 2A_4) \times L_2(7)).2 & (13:6 \times L_3(3)).2 & 13^{1+2}:(3 \times 4S_4) & L_2(71) \\ L_2(59) & L_2(41) & 41:40 & \end{array}$$

Let \mathcal{M}_1 denote this set of maximal subgroups of G and let x_1, \dots, x_t be a complete set of representatives of the conjugacy classes in G of elements of prime order. Let $\tau = (H_1, H_2, H_3)$ be a triple of subgroups with each $H_i \in \mathcal{M}_1$ and recall that

$$\widehat{Q}(G, \tau) = \sum_{i=1}^t |x_i^G| \cdot \left(\prod_{j=1}^3 \text{fpr}(x_i, G/H_j) \right).$$

We can use the stored fusion maps in GAP to compute $\text{fpr}(x_i, G/H)$ for all i and all $H \in \mathcal{M}_1$, whence

$$\widehat{Q}(G, \tau) \leq \sum_{i=1}^t a_i b_i^3, \quad (7)$$

where

$$a_i = |x_i^G|, \quad b_i = \max\{\text{fpr}(x_i, G/H) : H \in \mathcal{M}_1\}.$$

r	c_r		
2	11734592583376	23	90321336548400569373425664000000
3	500595349782528000	29	86565910978666325606400
5	9367743238695946498867200	31	134025209071820199715405824000000
7	3954417208796381184000	41	1680
11	3147561728201838023619379200000	47	88399605983540982791012352000000
13	44715330533129256960000	59	1740
17	61099727665094502811435008000000	71	2520
19	109336354769116478715199488000000		

TABLE 7. The values of c_r in the proof of Lemma 4.10

For example, if x_1 is a 2A-involution, then

$$a_1 = 97239461142009186000, \quad b_1 = \frac{56416310497}{467497409336582625}$$

(with $\text{fpr}(x_1, G/H) = b_1$ if $H = 2.\mathbb{B}$). It is entirely straightforward to check that the upper bound in (7) yields $\widehat{Q}(G, \tau) < 1$ and thus τ is regular.

So to complete the proof, we need to extend our analysis to triples containing one or more subgroups from the remaining 15 conjugacy classes of maximal subgroups H for which the fusion map from H -classes to G -classes is not available in [6].

Let r be a prime divisor of $|G|$ and define

$$i_r(H) = |\{x \in H : |x| = r\}|, \quad c_r = \min\{a_i b_i : |x_i| = r\}.$$

Notice that if

$$i_r(H) \leq c_r \tag{8}$$

for all r , then $\text{fpr}(x_i, G/H) \leq b_i$ for all i . So by our previous calculation, if \mathcal{S} is a collection of subgroups of G such that (8) holds for all r and all $H \in \mathcal{S}$, then every triple of subgroups in \mathcal{S} is regular. For the reader's convenience, the values of c_r are recorded in Table 7.

We claim that the inequality in (8) holds (for every prime r) when H is any one of the following maximal subgroups:

$$\begin{aligned} &(\text{L}_2(11) \times \text{L}_2(11)).4, 11^2:(5 \times 2A_5), \text{U}_3(4):4, \text{L}_2(29):2, 7^2:\text{SL}_2(7), \text{L}_2(19):2, \text{L}_2(13):2 \\ &3^{3+2+6+6}.\text{L}_3(3) \times \text{SD}_{16}, 3^{2+5+10}.\text{M}_{11} \times 2S_4, (3^2:2 \times \text{P}\Omega_8^+(3)).S_4 \\ &2^{3+6+12+18}.\text{L}_3(2) \times 3S_6, 2^{2+11+22}.\text{M}_{24} \times S_3 \end{aligned}$$

Indeed, in each case we can work with a permutation representation of H in the Web Atlas [49], which allows us to compute $i_r(H)$ and then verify the desired bound. For example, if $H = 2^{2+11+22}.\text{M}_{24} \times S_3$ then the Web Atlas provides a representation of H on 294912 points and we compute:

$$i_2(H) = 43521572863, \quad i_3(H) = 32114946867200, \quad i_5(H) = 68457483730944$$

$$i_7(H) = 1564742485278720, \quad i_{11}(H) = 23897885229711360, \quad i_{23}(H) = 182870773931704320,$$

noting that $i_r(H) = 0$ for all other primes. It is routine to check that (8) holds. Similarly, if $H = 2^{5+10+20}.\text{S}_3 \times \text{L}_5(2)$ or $2^{10+16}.\Omega_{10}^+(2)$, then one checks that the upper bound on $i_r(H)$ given in [16, Proposition 3.8] is sufficient.

Finally, let us assume $H = 3^8.\text{P}\Omega_8^-(3).2$. Here the Web Atlas provides a permutation representation of H on 805896 points and as before we can use this to compute $i_r(H)$ for every prime divisor r of $|G|$. In this way, we find that the inequality in (8) holds unless $r = 41$. More precisely, G has a unique class of elements of order 41 and we get

$$i_{41}(H) = 16245625881139200 > c_{41} = 1680.$$

However, if we now redefine b_i to be $i_{41}(H)/a_i$ (for the unique i such that $|x_i| = 41$) then it is easy to check that the inequality $\sum_i a_i b_i^3 < 1$ is still satisfied and we conclude that any maximal triple involving H is regular.

Putting all of this together, it follows that every maximal triple is regular and thus $R(G) = B(G) = 3$. \square

This completes the proof of Proposition 4.3. In particular, this concludes our proof of Theorem 2(i).

Remark 4.11. Let us briefly comment on the problem of determining all the non-regular maximal k -tuples for $G = \mathbb{B}, \mathbb{M}$ with $k = R(G) - 1$.

- (a) For $G = \mathbb{B}$ we have proved that $R(G) = B(G) = 4$, but we have not been able to classify all the non-regular triples. Let σ be a maximal triple and recall that we can compute $\widehat{Q}(G, \sigma)$ with the aid of the GAP Character Table Library [6]. In this way, we deduce that σ is non-regular only if it is one of the following:

$$(1, 1, 1), (1, 1, 2), (1, 1, 3), (1, 1, 4), (1, 1, 6), (1, 1, 7), (1, 2, 2),$$

where we are using the numbering in (6). We know that $(1, 1, 1)$ is non-regular by [16], and it is also easy to see that $(1, 1, 2)$ is non-regular by comparing orders. It remains an open problem to determine the regularity status of the following triples:

$$(1, 1, 3), (1, 1, 4), (1, 1, 6), (1, 1, 7), (1, 2, 2).$$

- (b) For the Monster $G = \mathbb{M}$ we have $R(G) = B(G) = 3$ and we have not attempted to determine the non-regular maximal pairs for G . From [16], we see that $(2.\mathbb{B}, 2.\mathbb{B})$ is the only non-regular conjugate maximal pair. In fact, [10, Theorem 3.1] implies that this is the only conjugate non-regular pair for G since $b(G, H) = 2$ for every nontrivial proper subgroup $H \neq 2.\mathbb{B}$. By comparing orders, we also note that $(2.\mathbb{B}, 2^{1+24}.\text{Co}_1)$ is non-regular.

4.2. Soluble subgroups. In this final section, we complete the proof of Theorem 2 by establishing the bound $R_{\text{sol}}(G) \leq 3$ for all almost simple sporadic groups. This generalises the main theorem of [10], where the weaker bound $B_{\text{sol}}(G) \leq 3$ is established. It also establishes a strong form of Conjecture 1(iii) in this setting. Our main result is the following.

Proposition 4.12. *We have $R_{\text{sol}}(G) \leq 3$ for every almost simple sporadic group G . More precisely,*

$$R_{\text{sol}}(G) = \begin{cases} 2 & \text{if } G \in \mathcal{A}_1 \\ 3 & \text{if } G \in \mathcal{A}_2 \\ 2 \text{ or } 3 & \text{if } G \in \mathcal{A}_3 \end{cases}$$

where

$$\begin{aligned} \mathcal{A}_1 &= \{J_1, J_3, J_3.2, \text{HS}, \text{Suz}, \text{Suz}.2, \text{McL}, \text{McL}.2, \text{Ru}, \text{He}, \text{He}.2, \text{Co}_3, \text{O}'\text{N}, \text{O}'\text{N}.2\} \\ \mathcal{A}_2 &= \{M_{11}, M_{12}, M_{12}.2, M_{22}, M_{22}.2, M_{23}, M_{24}, J_2, J_2.2, \text{HS}.2, \text{Co}_2, \text{Fi}_{22}, \text{Fi}_{22}.2, \text{Fi}_{23}\} \\ \mathcal{A}_3 &= \{\text{Co}_1, \text{HN}, \text{HN}.2, J_4, \text{Ly}, \text{Th}, \text{Fi}'_{24}, \text{Fi}_{24}, \mathbb{B}, \mathbb{M}\} \end{aligned}$$

Remark 4.13. Let us record some comments on the statement of Proposition 4.12.

- (a) Clearly, we have $R_{\text{sol}}(G) \leq R(G)$, so the main bound $R_{\text{sol}}(G) \leq 3$ follows immediately from Proposition 4.3 when G is one of the following:

$$J_1, J_3, J_3.2, \text{O}'\text{N}, \text{O}'\text{N}.2, \text{HN}, \text{HN}.2, J_4, \text{Ly}, \text{Th}, \mathbb{M}.$$

- (b) We can determine all the non-regular soluble pairs for the following groups

$$M_{11}, M_{12}, M_{12}.2, M_{22}, M_{22}.2, M_{23}, M_{24}, J_2, J_2.2, \text{HS}.2 \quad (9)$$

In particular, if $G = M_{23}$, then (H, H) is the only non-regular soluble pair, up to conjugacy, where

$$H = 2^4:(3 \times A_4):2 < 2^4:(3 \times A_5):2 < G$$

G	H	K	Comments
M_{11}	$U_3(2):2$	$U_3(2)$	
M_{12}	$AGL_2(3)$	$ASL_2(3)$	
$M_{12}.2$	$AGL_2(3)$	$4^2:D_{12}.2$	$H < M_{12} < G, K < G$
M_{22}	$2^4:(3^2:4)$	$2^4:S_4$	$H < 2^4:A_6 < G, K < 2^4:A_6 < G$
$M_{22}.2$	$2^4:(S_3 \wr S_2)$	$2^4:(S_4 \times S_2)$	$H < 2^4:S_6 < G, K < 2^4:S_6 < G$
M_{24}	$2^6:3.(S_3 \wr S_2)$	$2^6:(7:3 \times S_3)$	$H < 2^6:3.S_6 < G, K < 2^6:(L_3(2) \times S_3) < G$
J_2	$2^{2+4}:(3 \times S_3)$	$2^{2+4}:3^2$	
$J_2.2$	$2^{2+4}:(3 \times S_3).2$	$2^{2+4}:(3 \times S_3)$	

TABLE 8. The subgroups H and K in Remark 4.13(b)

is a second maximal subgroup of G . Similarly, $G = HS.2$ has a unique non-regular soluble pair, up to ordering and conjugacy, namely (H, K) , where

$$H = 2^{1+6}.S_4 < 2^{1+6}.S_5 < G$$

$$K = 4^3:(2 \times 7:3) < 4^3:(2 \times L_3(2)) < G$$

are both second maximal subgroups. In Table 8, for each of the remaining groups in (9), we record a non-regular conjugate pair (H, H) with $|H|$ maximal, and also a non-regular non-conjugate pair (H, K) with $|H||K|$ maximal (in every case, K is a second maximal subgroup of G , and H is either maximal or second maximal).

- (c) For the remaining groups in \mathcal{A}_2 , namely Co_2 , Fi_{22} , $Fi_{22}.2$ and Fi_{23} , a non-regular conjugate soluble pair (H, H) is recorded in [10, Table 1]. But in these cases we have not classified all the non-regular soluble pairs. By inspecting [10, Table 2], we observe that (H, K) is a non-regular non-conjugate pair for $G = Fi_{22}.2$, where the first component $H = 3^{1+6}.2^{3+4}.3^2:2.2$ is maximal and $K = 3^{1+6}.2^{3+4}.3^2:2 < H$ is an index-two subgroup. Similarly, if $G = Fi_{23}$ then (H, K) is non-regular, where $H = 3^{1+8}.2^{1+6}.3^{1+2}.2S_4$ is maximal and K is the unique index-two subgroup of H .
- (d) We have not been able to determine $R_{sol}(G)$ precisely for the groups in \mathcal{A}_3 . By [10, Theorem 2], none of these groups, with the possible exception of the Baby Monster \mathbb{B} , has a non-regular conjugate soluble pair. Determining the existence (or otherwise) of such a pair for \mathbb{B} remains an open problem.
- (e) We refer the reader to the end of Section 4.2 for some brief comments regarding the regularity of nilpotent pairs (H, K) in almost simple sporadic groups, as well as the pairs (H, K) with H nilpotent and K soluble.

Let G be an almost simple sporadic group with socle T . In order to prove Proposition 4.12 we will adopt a computational approach, working with MAGMA (version 2.28-4) [5] and GAP (version 4.11.1). It will be convenient to divide up the possibilities for T as follows:

$$\mathcal{B}_1 = \{M_{11}, M_{12}, M_{22}, M_{23}, M_{24}, J_1, J_2, J_3, HS, McL, He, Suz, Ru, Co_3, O'N\}$$

$$\mathcal{B}_2 = \{Co_2, Fi_{22}, Fi_{23}, HN, J_4, Ly, Th, \mathbb{M}\}$$

$$\mathcal{B}_3 = \{Co_1, Fi'_{24}\}$$

$$\mathcal{B}_4 = \{\mathbb{B}\}$$

Before we begin the proof, it will be helpful to introduce the following notation. For a positive integer k , let $\mathcal{C}_k(G)$ be the set of subgroups H of G with the property that there exists a chain of subgroups

$$H = H_0 < H_1 < H_2 < \cdots < H_{k-1} < H_k = G$$

where H_1 is insoluble and each H_i is a maximal subgroup of H_{i+1} . For instance, $\mathcal{C}_1(G)$ is the set of maximal subgroups of G . In addition, let $\mathcal{D}_k(G)$ be the set of soluble subgroups in $\mathcal{C}_k(G)$, and let $\mathcal{P}(G)$ be the set of all non-Sylow p -subgroups of G , ranging over all prime divisors p of $|G|$.

Definition 4.14. Let $H \leq G$ be a subgroup and let d be a positive integer. If H is soluble, we define $\mathcal{S}_d(H) = \{H\}$. Otherwise, we take $\mathcal{S}_d(H)$ to be a set of representatives of the G -classes of subgroups in the set

$$\left(\mathcal{C}_d(H) \cup \bigcup_{k=1}^{d-1} \mathcal{D}_k(H) \right) \setminus \mathcal{P}(H)$$

For example, if $G = H = M_{11}$, then $\mathcal{S}_2(H)$ contains the soluble maximal subgroups $U_3(2):2$ and $2S_4$, together with a representative of each conjugacy class of maximal subgroups of $M_{10} = A_6.2, L_2(11)$ and S_5 .

The key property here is that for any positive integer d , every soluble subgroup of H is G -conjugate to a subgroup of at least one group in $\mathcal{S}_d(H)$. It follows that $R_{\text{sol}}(G) \leq 3$ if there exist positive integers d_j such that every triple $\tau = (H_1, H_2, H_3)$ is regular, where the components of τ range over the subgroups in

$$\bigcup_{j=1}^t \mathcal{S}_{d_j}(M_j)$$

with respect to a complete set of representatives $\{M_1, \dots, M_t\}$ of the conjugacy classes of maximal subgroups of G .

Lemma 4.15. *The conclusion to Proposition 4.12 holds for the groups with $T \in \mathcal{B}_1$.*

Proof. Let G be an almost simple sporadic group with socle $T \in \mathcal{B}_1$. Working with MAGMA, we first use the function `AutomorphismGroupSimpleGroup` to construct a faithful primitive permutation representation of G of minimal degree. We then use `SolubleSubgroups` to obtain a complete set of representatives of the conjugacy classes of soluble subgroups of G . Let $\mathcal{S}(G)$ be the set obtained by removing all the non-Sylow p -subgroups, ranging over all the prime divisors p of $|G|$. We then use random search to determine a list of candidate non-regular soluble pairs (H_1, H_2) , where $|H_1| \geq |H_2|$ and $H_i \in \mathcal{S}(G)$ for $i = 1, 2$.

Clearly, if random search reveals that every pair (H_1, H_2) with $H_i \in \mathcal{S}(G)$ is regular, then $R_{\text{sol}}(G) = 2$. So let us assume that there is at least one such pair (H_1, H_2) for which our random search is inconclusive. Then as in the proof of Proposition 4.3, we can determine the regularity status of (H_1, H_2) by computing the orbits of H_2 on G/H_1 via the function `CosetAction`. In this way, we obtain the complete list of non-regular soluble pairs, up to ordering and conjugacy, and we can easily extract the information highlighted in part (b) of Remark 4.13. Of course, if we find that there are no such pairs, then $R_{\text{sol}}(G) = 2$.

To complete the argument, let us assume G has at least one non-regular soluble pair. In order to show that every soluble triple is regular, we consider each non-regular soluble pair (H_1, H_2) in turn, and we construct the triples $\tau = (H_1, H_2, H_3)$, where H_3 runs through the set $\mathcal{S}(G)$. Using random search, it is straightforward to verify that τ is regular and we conclude that $R_{\text{sol}}(G) = 3$ in each of these cases. \square

Lemma 4.16. *The conclusion to Proposition 4.12 holds for the groups with $T \in \mathcal{B}_2$.*

Proof. If $T \in \{\text{HN}, J_4, \text{Ly}, \text{Th}, \text{M}\}$ then $R(G) = 3$ by Proposition 4.3 and thus $R_{\text{sol}}(G) \leq 3$, as required. For the remainder, let us assume $T \in \{\text{Co}_2, \text{Fi}_{22}, \text{Fi}_{23}\}$. In these cases, the main theorem of [10] gives $B_{\text{sol}}(G) = 3$, so we just need to show that every soluble triple is regular in order to conclude that $R_{\text{sol}}(G) = 3$. To do this, we need to modify the computational approach adopted in the proof of Lemma 4.15 since the MAGMA function `SolubleSubgroups` is ineffective for the groups we are working with here. We proceed as follows.

For now, let us assume $G \neq \text{Fi}_{22}.2$, so $G = T$. As before, we begin by constructing G as a permutation group and we use the MAGMA function `MaximalSubgroups` to obtain a complete set of representatives of the conjugacy classes of maximal subgroups of G . Then for each maximal subgroup H , we construct the set of subgroups $\mathcal{S}_3(H)$ as defined in Definition 4.14. In this way, we obtain a set of subgroups with the property that every soluble subgroup of G

is contained in a conjugate of at least one member of the set. We can then use random search to show that every triple of the form (J_1, J_2, J_3) is regular, where each J_i is contained in $\mathcal{S}_3(H_i)$ for some maximal subgroup H_i of G , and this allows us to conclude that $R_{\text{sol}}(G) \leq 3$.

In practice, to improve the efficiency of the latter computation, we first run through triples of the form (J_1, H_2, H_3) , where $J_1 \in \mathcal{S}_3(H_1)$ and the H_i are representatives of the classes of maximal subgroups of G . Given H_2 and H_3 , if random search is inconclusive for at least one triple of the form (J_1, H_2, H_3) , then we run through the set of triples $\tau = (J_1, J_2, H_3)$ with $J_i \in \mathcal{S}_3(H_i)$ for $i = 1, 2$, and in every case we find that τ is regular.

Finally, suppose $G = \text{Fi}_{22}.2$. We know that $R_{\text{sol}}(T) = 3$ by the above argument, so Lemma 2.11 implies that every soluble triple of the form (H, K, L) with $H < T$ is regular. So we can proceed as above, using the additional fact that we can discard any triples with a component contained in T . \square

Lemma 4.17. *The conclusion to Proposition 4.12 holds for the groups with $T \in \mathcal{B}_3$.*

Proof. First assume $G = \text{Fi}'_{24}$. Here we proceed as in the proof of Proposition 4.3, using the information in the GAP Character Table Library [6] to determine all the maximal triples σ with $\widehat{Q}(G, \sigma) \geq 1$. It turns out that this inequality holds only if at least one of the component subgroups is Fi_{23} . But we know that $R_{\text{sol}}(\text{Fi}_{23}) = 3$ by Lemma 4.16, so Lemma 2.11 implies that every soluble triple is regular and thus $R_{\text{sol}}(G) \leq 3$.

Next assume $G = \text{Co}_1$. Here we observe that the largest maximal subgroup of G is Co_2 , and we have $R_{\text{sol}}(\text{Co}_2) = 3$ by Lemma 4.16. Similarly, $R_{\text{sol}}(\text{Co}_3) = 2$ by Lemma 4.15. Therefore, Lemma 2.11 implies that a soluble triple is regular if it has a component contained in one of the maximal subgroups Co_2 or Co_3 of G . We can now use GAP, as in the previous case, to determine all the maximal triples $\sigma = (H_1, H_2, H_3)$, up to ordering and conjugacy, such that $\widehat{Q}(G, \sigma) \geq 1$ and $H_i \neq \text{Co}_2, \text{Co}_3$ for all i . We find that there are just 12 such triples, up to conjugacy and reordering, which only involve subgroups from 6 distinct conjugacy classes of maximal subgroups of G .

At this point, we switch to MAGMA and we construct G and each of the 6 relevant maximal subgroups inside S_n , where $n = |G : \text{Co}_2| = 98280$. Using random search, we find that 10 of the 12 relevant maximal triples are regular and it just remains to consider the maximal triples (H_1, H_2, H_3) , where

$$H_1 = H_2 = 3.\text{Suz}:2, \quad H_3 = 3.\text{Suz}:2 \text{ or } 2^{11}:\text{M}_{24}.$$

We now complete the argument by proceeding as in the proof of Lemma 4.16, working with the subgroup collections $\mathcal{S}_3(H_i)$ and random search to show that each triple of the form (J_1, H_2, H_3) with $J_1 \in \mathcal{S}_3(H_1)$ is regular. In this way, we conclude that every soluble triple of subgroups of G is regular and thus $R_{\text{sol}}(G) \leq 3$.

Finally, let us assume $G = \text{Fi}_{24}$. Since we have already established the bound $R_{\text{sol}}(T) \leq 3$, Lemma 2.11 implies that every soluble triple containing a subgroup of T is regular.

We begin by using GAP to determine all the core-free maximal triples σ with $\widehat{Q}(G, \sigma) \geq 1$. Switching to MAGMA, we then use the Web Atlas [49] to construct G and each relevant maximal subgroup H_i as subgroups of S_n , where $n = 306936$. By applying random search, we can then reduce the candidate non-regular core-free maximal triples (H_1, H_2, H_3) to a specific list of 18 possibilities (up to conjugacy and reordering), all of which include the component $H_1 = \text{Fi}_{23} \times 2$. Next we construct the set $\mathcal{S}_3(H_1)$ defined in Definition 4.14. By removing any subgroups that are contained in T , we obtain a subset $\mathcal{S}_3(H_1)'$ of size 103. Then by applying random search, we can show that every triple of the form (J_1, H_2, H_3) is regular, where $J_i \in \mathcal{S}_3(H_1)'$, unless $H_2 = H_1$ and H_3 is one of the following:

$$\text{Fi}_{23} \times 2, \quad (2 \times 2.\text{Fi}_{22}):2, \quad \text{S}_3 \times \text{P}\Omega_8^+(3):\text{S}_3, \quad \text{O}_{10}^-(2).$$

Finally, we handle these cases by showing that every triple of the form (J_1, J_2, H_3) is regular (via random search), where $J_i \in \mathcal{S}_3(H_1)'$ for $i = 1, 2$. \square

To complete the proof of Proposition 4.12, we may assume $G = \mathbb{B}$ is the Baby Monster.

Lemma 4.18. *The conclusion to Proposition 4.12 holds for the groups with $T \in \mathcal{B}_4$.*

Proof. Here $G = \mathbb{B}$ and we begin by considering a maximal triple σ . As noted in the proof of Lemma 4.10, we can use the GAP Character Table Library [6] to show that $\widehat{Q}(G, \sigma) < 1$ unless σ is of the form $(1, 1, i)$ or $(1, 2, 2)$, where $i \in \{1, 2, 3, 4, 6, 7\}$ and the relevant subgroups are labelled as in (6). We immediately deduce that if τ is a soluble triple and none of the components are contained in a maximal subgroup $2.^2E_6(2):2$, then τ is regular.

For the remainder of the proof, set

$$L = 2.^2E_6(2):2, \quad Z = Z(L) = \langle z \rangle, \quad \bar{L} = L/Z = {}^2E_6(2):2.$$

As in [10, Section 4.1], write $\bar{L}' = {}^2E_6(2) = (X_\sigma)'$, where $X = E_6$ is a simple algebraic group of adjoint type over an algebraically closed field of even characteristic and σ is a Steinberg endomorphism of X such that

$$X_\sigma = \{x \in X : x^\sigma = x\} = \text{Inndiag}({}^2E_6(2)) = {}^2E_6(2):3.$$

Let \mathcal{M}_0 be a set of representatives of the conjugacy classes of maximal subgroups of G , excluding L , and define

$$\mathcal{M} = \mathcal{M}_1 \cup \mathcal{M}_2,$$

where

$$\begin{aligned} \mathcal{M}_1 &= \{\text{maximal subgroups of } L, \text{ other than } 2.^2E_6(2)\} \\ \mathcal{M}_2 &= \{\text{maximal subgroups of } 2.^2E_6(2) < L\} \end{aligned}$$

Note that every soluble subgroup of L is contained in a subgroup $K \in \mathcal{M}$. Then in view of the above GAP calculation, it suffices to prove the following claim:

Claim. *Let $\sigma = (H_1, H_2, H_3)$ be a triple of subgroups of G , where $H_1 \in \mathcal{M}$ and $H_2, H_3 \in \mathcal{M}_0 \cup \mathcal{M}$. Then $\widehat{Q}(G, \sigma) < 1$.*

To prove the claim, we proceed as in the proof of Lemma 4.10, where we showed that $R(\mathbb{M}) = 3$. First observe that we can use the GAP Character Table Library [6] to compute $\text{fpr}(x, G/H)$ for every maximal subgroup H and every element $x \in G$. Let x_1, \dots, x_t be a complete set of representatives of the conjugacy classes in G of elements of prime order and set $a_i = |x_i^G|$ and

$$b_i = \max\{\text{fpr}(x_i, G/H) : H \in \mathcal{M}_0\} \quad (10)$$

for all $1 \leq i \leq t$. It follows immediately that the bound in (7) holds for every maximal triple $\tau = (H_1, H_2, H_3)$ with $H_i \in \mathcal{M}_0$ for all i . Moreover, we check that this upper bound yields $\widehat{Q}(G, \tau) < 1$ and thus τ is regular. So in order to prove the claim, it suffices to show that

$$\text{fpr}(x_i, G/H) \leq b_i \quad (11)$$

for all $1 \leq i \leq t$ and all $H \in \mathcal{M}$. For the reader's convenience, the values of b_i are recorded in Table 9, with respect to the standard labelling of the conjugacy classes in G (see [22]).

Fix a subgroup $H \in \mathcal{M}_2$, so H is a maximal subgroup of $J = 2.^2E_6(2)$. Then $Z \leq H$ and thus $\bar{H} = H/Z$ is a maximal subgroup of the simple group $\bar{J} = J/Z = {}^2E_6(2)$. The possibilities for \bar{H} (up to conjugacy) have been determined by Wilson [51], which confirms that the list of maximal subgroups presented in the ATLAS [22] is complete (also see [23]). We observe that $\bar{H} = \bar{K} \cap \bar{J}$ for some core-free maximal subgroup $\bar{K} = K/Z$ of $\bar{L} = L/Z = {}^2E_6(2):2$, whence $H \leq K$ and K is contained in \mathcal{M}_1 . Therefore, we only need to prove that (11) holds for each subgroup $H \in \mathcal{M}_1$.

Suppose $H \in \mathcal{M}_1$, so $H \neq 2.^2E_6(2)$ is a maximal subgroup of $L = 2.^2E_6(2):2$. Then $Z \leq H$ and $\bar{H} = H/Z$ is a maximal subgroup of the almost simple group \bar{L} . As above, the

i	x_i	b_i
1	2A	793/2950425
2	2B	269689951/11707448673375
3	2C	13/33634845
4	2D	303281/780496578225
5	3A	11/105887475
6	3B	2/5353200125
7	5A	11/22299902235
8	5B	2/93659589387
9	7A	1/37166503725
10	11A	1/780496578225
11	13A	1/126996316160000
12	17A	1/253992632320000
13	19A	1/22892381208576000
14	23A	1/11707448673375
15	23B	1/11707448673375
16	31A	1/45784762417152000
17	31B	1/45784762417152000
18	47A	1/3843461129719173164826624000000
19	47B	1/3843461129719173164826624000000

TABLE 9. The b_i values in (10)

possibilities for \bar{H} have been determined by Wilson [51] and we deduce that \bar{H} is one of the following:

- Parabolic: $P_{1,6}, P_2, P_{3,5}, P_4$
- Algebraic: $O_{10}^-(2), S_3 \times U_6(2):2, S_3 \times \Omega_8^+(2):S_3, U_3(8):6, (L_3(2) \times L_3(4):2).2,$
 $3^{1+6}.2^{3+6}.3^2.2^2, U_3(2):2 \times G_2(2), F_4(2) \times 2$
- Almost simple: $SO_7(3), Fi_{22}:2$

Here we adopt the standard notation for the maximal parabolic subgroups of \bar{L} , which corresponds to the usual labelling of the nodes of the Dynkin diagram of type E_6 . The algebraic subgroups are of the form $N_{\bar{L}}(Y_{\sigma})$, where Y is a positive dimensional non-parabolic σ -stable closed subgroup of the ambient algebraic group X . For example, if $\bar{H} = 3^{1+6}.2^{3+6}.3^2.2^2$ then the connected component of Y is of type A_2^3 and we will refer to $SU_3(2)^3$ as the *type* of \bar{H} . We extend this usage of type to the other algebraic subgroups, which provides an approximate description of the given subgroup's structure.

Let $\pi(\bar{H})$ be the set of prime divisors of $|\bar{H}|$ and recall that our goal is to verify the bound in (11) for all i . Let $i_r(H)$ be the number of elements of order r in H and observe that

$$i_2(H) = 2i_2(\bar{H}) + 1, \quad i_r(H) = i_r(\bar{H})$$

for every odd prime r . We now consider each possibility for \bar{H} in turn.

Case (a). $\bar{H} = (L_3(2) \times L_3(4):2).2$

Let $x \in G$ be an element of prime order $r \in \pi(\bar{H}) = \{2, 3, 5, 7\}$ and note that $|H| = 2|\bar{H}| = 27095040$. It is easy to check that the trivial bound $|x^G \cap H| \leq |H|$ is sufficient unless $x \in 2A$. Since $\bar{H} \leq A$, where $A = \text{Aut}(L_3(2) \times L_3(4))$, we deduce that $i_2(\bar{H}) \leq i_2(A) = 98199$ and thus

$$|x^G \cap H| \leq i_2(H) \leq 196399$$

for $x \in 2A$. This bound is sufficient.

Case (b). $\bar{H} = 3^{1+6}.2^{3+6}.3^2.2^2$

Here $\pi(\bar{H}) = \{2, 3\}$ and once again we find that the trivial bound $|x^G \cap H| \leq |H|$ is good enough unless $x \in 2A$. So let us assume $x \in 2A$. Then by inspecting the proof of [18, Lemma

4.12], noting that $i_2(\mathrm{SU}_3(2)) = 9$, we deduce that $i_2(\bar{H}) \leq \alpha + \beta$, where

$$\alpha = \binom{3}{1}9 + \binom{3}{1}9^2 + 9^3 + \binom{3}{2}|\mathrm{SU}_3(2)| \cdot \frac{|\mathrm{SU}_3(2)|}{|\mathrm{SL}_2(2)|}$$

$$\beta = 3(1+9)|\mathrm{SU}_3(2)| + \left(\frac{|\mathrm{SU}_3(2)|}{|\mathrm{SL}_2(2)|} \right)^3$$

This yields $|x^G \cap H| \leq i_2(H) \leq 154927$ and the desired bound follows.

Case (c). $\bar{H} = S_3 \times \mathrm{U}_6(2):2$, $S_3 \times \Omega_8^+(2):S_3$, $\mathrm{U}_3(8):6$ or $\mathrm{U}_3(2):2 \times G_2(2)$

In each of these cases, with the aid of MAGMA, it is easy to compute $i_r(H)$ for all $r \in \pi(H)$ and we find that the bound $|x^G \cap H| \leq i_r(H)$ is sufficient unless $\bar{H} = S_3 \times \mathrm{U}_6(2):2$ and $x \in 2A$. In the latter case, the proof of [10, Lemma 4.6] gives $|x^G \cap H| \leq 7033$ and this bound is good enough.

Case (d). $\bar{H} = \mathrm{O}_{10}^-(2)$, $\mathrm{SO}_7(3)$, $\mathrm{Fi}_{22}:2$ or $F_4(2) \times 2$

Let $x \in G$ be an element of prime order r . If $r = 2$ then $|x^G \cap H|$ is recorded in [10, Table 5] and the desired bound quickly follows (see Remark 4.19). Now assume r is odd. Here we can compute $i_r(H)$ from the character table of \bar{H} , which is available in [6], and in every case it is easy to check that the bound $|x^G \cap H| \leq i_r(H)$ is sufficient.

Case (e). $\bar{H} = P_{1,6}$, P_2 , $P_{3,5}$ or P_4

To complete the proof, we may assume \bar{H} is a maximal parabolic subgroup of \bar{L} . Note that the precise structure of \bar{H} is recorded in [10, Table 6]. In addition, let us observe that $\pi(H) = \{2, 3, 5, 7\} \cup \mathcal{P}$, where $\mathcal{P} = \{17\}$ for P_1 , $\mathcal{P} = \{11\}$ for $P_{1,6}$, and \mathcal{P} is empty for $P_{3,5}$ and P_4 . Let $x \in G$ be an element of order $r \in \pi(H)$.

If $r = 2$ or 3 , then $|x^G \cap H|$ is recorded in [10, Tables 5 and 6] and it is easy to check that (11) holds for all relevant i . Now assume $r \in \{5, 7\} \cup \mathcal{P}$. Here one can check that the trivial bound

$$|x^G \cap H| \leq |H| \leq 2|P_2| = 2^{23}|\mathrm{U}_6(2)|$$

is sufficient for $r \geq 7$, so we may assume $r = 5$ and thus $x \in 5A$ or $5B$. Now L has a unique conjugacy class of elements of order 5 and the fusion map from L -classes to G -classes shows that this class is contained in the G -class labelled $5A$. So we may assume $x \in 5A$ and thus

$$|x^G \cap H| = i_5(H) = i_5(\bar{H}).$$

In addition, let us observe that if $\bar{H} = P_{3,5}$ or P_4 , then $|H| \leq 2|P_4| = 2^{32}3|\mathrm{L}_3(4)|$ and the crude bound $|x^G \cap H| \leq 2|P_4|$ is good enough. Therefore, we are free to assume that $\bar{H} = P_{1,6}$ or P_2 .

Let $x \in H$ be an element of order 5 and let \bar{x} be the image of x in $\bar{H} = H/Z$. Then

$$\mathrm{fpr}(\bar{x}, \bar{L}/\bar{H}) = \frac{i_5(\bar{H})}{|\bar{x}^{\bar{L}}|}$$

and by applying [35, Theorem 2], noting that $\bar{x} \in \bar{L}$ is semisimple, we deduce that

$$|x^G \cap H| = i_5(\bar{H}) \leq |\bar{x}^{\bar{L}}| \cdot \frac{1}{3 \cdot 2^6}.$$

Now

$$|\bar{x}^{\bar{L}}| = 759250790513639424, \quad |x^G| = 9367743238695946498867200$$

and we conclude that

$$\mathrm{fpr}(x, G/H) \leq \frac{3954431200591872}{9367743238695946498867200} < \frac{11}{22299902235} = b_7,$$

as required. \square

This completes the proof of Theorem 2.

G	H	K
M_{11}	$U_3(2):2$	$U_3(2):2, 2S_4$
M_{12}	$3^2:2S_4$	$3^2:2S_4, 2^{1+4}:S_3, 4^2:D_{12}$
	$2^{1+4}:S_3$	$2^{1+4}:S_3, 4^2:D_{12}$
	$4^2:D_{12}$	$4^2:D_{12}$
$M_{12}.2$	$2^{1+4}:S_3.2$	$2^{1+4}:S_3.2, 4^2:D_{12}.2, 3^{1+2}:D_8$
	$4^2:D_{12}.2$	$4^2:D_{12}.2$
	$3^{1+2}:D_8$	$3^{1+2}:D_8$
J_2	$2^{2+4}:(3 \times S_3)$	$2^{2+4}:(3 \times S_3), 5^2:D_{12}$
$J_2.2$	$2^{2+4}:(3 \times S_3).2$	$2^{2+4}:(3 \times S_3).2, 5^2:(4 \times S_3)$
Fi_{22}	$3^{1+6}.2^{3+4}.3^2.2$	$3^{1+6}.2^{3+4}.3^2.2$
$Fi_{22}.2$	$3^{1+6}.2^{3+4}.3^2.2.2$	$3^{1+6}.2^{3+4}.3^2.2.2$
Fi_{23}	$3^{1+8}.2^{1+6}.3^{1+2}.2S_4$	$3^{1+8}.2^{1+6}.3^{1+2}.2S_4$

TABLE 10. The pairs (H, K) in Proposition 4.20

Remark 4.19. In the proof of Lemma 4.18 we appealed to the information in [10, Table 5]. Here we take the opportunity to correct three inaccuracies in this table:

- (a) For $\bar{H} = O_{10}^-(2)$ we have $|2D \cap H| = 79942500$, rather than 79943000.
- (b) And for $\bar{H} = F_4(2) \times 2$, the stated sizes of $2A \cap H$ and $2C \cap H$ are out by 1; the correct values are 139232 and 355284576, respectively.

We finish by presenting Proposition 4.20 below, which gives the exact value of $R_{\text{sol max}}(G)$ for every almost simple group G with a soluble maximal subgroup. By inspecting [50], for example, it is easy to see that G has such a subgroup unless $G = M_{22}, M_{22}.2, M_{24}$ or HS.

Proposition 4.20. *Let G be an almost simple sporadic group with socle T and assume G has a soluble maximal subgroup. Then*

$$R_{\text{sol max}}(G) = \begin{cases} 3 & \text{if } T = M_{11}, M_{12}, J_2, Fi_{22} \text{ or } Fi_{23} \\ 2 & \text{otherwise.} \end{cases}$$

Moreover, if $R_{\text{sol max}}(G) = 3$ then every soluble maximal non-regular pair (H, K) is recorded in Table 10, up to conjugacy and ordering.

Proof. As noted above, the assumption that G has a soluble maximal subgroup means that $G \notin \{M_{22}, M_{22}.2, M_{24}, HS\}$. In addition, if

$$G \in \{M_{23}, HS.2, McL, Suz, Suz.2, Ru, Co_3, Co_2, O'N, Fi_{22}, Fi_{22}.2, Fi_{23}, Fi'_{24}\},$$

then G has a unique class of soluble maximal subgroups, so every soluble maximal pair is conjugate and the result follows from [16].

Next assume $G \in \{M_{11}, M_{12}, M_{12}.2, J_2, J_2.2\}$ is one of the remaining groups appearing in Table 10. In each of these cases, an entirely straightforward MAGMA computation shows that a soluble maximal pair (H, K) is non-regular if and only if it is one of the cases recorded in the table (up to conjugacy and ordering).

In each of the remaining cases, we can use the GAP Character Table Library [6] to compute $\widehat{Q}(G, \tau)$ precisely for every soluble maximal pair $\tau = (H, K)$. In this way, we deduce that $\widehat{Q}(G, \tau) < 1$, which implies that τ is regular, unless $G = \text{He}.2$ and $H = K = 2^{4+4}:(S_3 \times S_3).2$, or $G = J_3.2$ and $H = K = 2^{2+4}:(S_3 \times S_3)$. In both cases, the main theorem of [16] implies that τ is regular. \square

Remark 4.21. The group M_{12} has two distinct classes of soluble maximal subgroups isomorphic to $3^2:2S_4$; in Table 10, we write $3^2:2S_4$ for a representative of either one of these classes.

Remark 4.22. Zenkov's main theorem in [55] states that every nilpotent pair of subgroups in an almost simple sporadic group is regular. Here we briefly comment on the existence (or otherwise) of non-regular pairs (H, K) , where H is nilpotent and K is soluble.

- (a) Clearly, if $R_{\text{sol}}(G) = 2$, then every nilpotent-soluble pair is regular, so we may assume $G \in \mathcal{A}_2 \cup \mathcal{A}_3$ as defined in the statement of Proposition 4.12.
- (b) If $G = M_{22}.2$ then there are precisely 7 non-regular pairs (H, K) , up to ordering and conjugation, where H is nilpotent and K is soluble. For example, we can take H to be a Sylow 2-subgroup and K to be the following second maximal subgroup

$$K = 2^4:(S_3 \wr S_2) < 2^4:S_6 < G.$$

For each such pair (H, K) we find that K has Fitting length 3, which is the maximal Fitting length of all soluble subgroups of G .

- (c) Similarly, $G = J_2.2$ has a unique non-regular pair (H, K) of this form. Here H is a Sylow 2-subgroup and $K = 2^{2+4}:(3 \times S_3).2$ is a maximal subgroup of G . Once again, the Fitting length of K is 3 and one checks this is the maximal Fitting length of all soluble subgroups of G .

We have checked computationally that the examples in (b) and (c) are the only non-regular nilpotent-soluble pairs associated with any of the groups in \mathcal{A}_2 . We are not aware of any additional examples for the groups in \mathcal{A}_3 .

REFERENCES

- [1] M. Anagnostopoulou-Merkouri and T.C. Burness, *Computations concerning the regularity number of almost simple groups*, 2024. <https://seis.bristol.ac.uk/~tb13602/regcomp.pdf>
- [2] L. Babai, A. Goodman and L. Pyber, *Groups without faithful transitive permutation representations of small degree*, J. Algebra **195** (1997), 1–29.
- [3] A.A. Baikalov, *Intersection of conjugate solvable subgroups in symmetric groups*, Algebra Logic **56** (2017), 87–97.
- [4] R.F. Bailey and P.J. Cameron, *Base size, metric dimension and other invariants of groups and graphs*, Bull. Lond. Math. Soc. **43** (2011), 209–242.
- [5] W. Bosma, J. Cannon and C. Playoust, *The MAGMA algebra system I: The user language*, J. Symb. Comput. **24** (1997), 235–265.
- [6] T. Breuer, *The GAP Character Table Library, Version 1.3.3*, GAP package, <http://www.math.rwth-aachen.de/~Thomas.Breuer/ctbllib>, 2022.
- [7] T.C. Burness, *On base sizes for actions of finite classical groups*, J. London Math. Soc. **75** (2007), 545–562.
- [8] T.C. Burness, *Simple groups, fixed point ratios and applications*, in Local representation theory and simple groups, 267–322, EMS Ser. Lect. Math., Eur. Math. Soc., Zürich, 2018.
- [9] T.C. Burness, *Base sizes for primitive groups with soluble stabilisers*, Algebra and Number Theory **15(7)** (2021), 1755–1807.
- [10] T.C. Burness, *On soluble subgroups of sporadic groups*, Israel J. Math. **254** (2023), 313–340.
- [11] T.C. Burness, M. Garonzi and A. Lucchini, *On the minimal dimension of a finite simple group*, J. Combin. Theory Ser. A **171** (2020), 105175, 32pp.
- [12] T.C. Burness, M. Garonzi and A. Lucchini, *Finite groups, minimal bases and the intersection number*, Trans. London Math. Soc. **9** (2022), 20–55.
- [13] T.C. Burness and R.M. Guralnick, *Strongly base-two groups*, Vietnam J. Math. **51** (2023), 657–683.
- [14] T.C. Burness, R.M. Guralnick and J. Saxl, *On base sizes for symmetric groups*, Bull. Lond. Math. Soc. **43** (2011), 386–391.
- [15] T.C. Burness, M.W. Liebeck and A. Shalev, *Base sizes for simple groups and a conjecture of Cameron*, Proc. Lond. Math. Soc. **98** (2009), 116–162.
- [16] T.C. Burness, E.A. O’Brien and R.A. Wilson, *Base sizes for sporadic simple groups*, Israel J. Math. **177** (2010), 307–333.
- [17] T.C. Burness and Á. Seress, *On Pyber’s base size conjecture*, Trans. Amer. Math. Soc. **367** (2015), 5633–5651.
- [18] T.C. Burness and A.R. Thomas, *The classification of extremely primitive groups*, Int. Math. Res. Not. IMRN (2022), 10148–10248.
- [19] P.J. Cameron, *Some measures of finite groups related to permutation bases*, preprint (arXiv:1408.0968), 2014.
- [20] P.J. Cameron, *Permutation Groups*, London Math. Soc. Student Texts **45**, Cambridge University Press, 1999.
- [21] P.J. Cameron and W.M. Kantor, *Random permutations: some group-theoretic aspects*, Combin. Probab. Comput. **2** (1993), 257–262.

- [22] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker and R.A. Wilson, *Atlas of finite groups*, Oxford University Press, 1985.
- [23] D.A. Craven, *The maximal subgroups of the exceptional groups $F_4(q)$, $E_6(q)$ and ${}^2E_6(q)$ and related almost simple groups*, *Invent. Math.* **234** (2023), 637–719.
- [24] C. del Valle and C.M. Roney-Dougal, *The base size of the symmetric group acting on subsets*, *Algebr. Comb.*, to appear.
- [25] H. Dietrich, M. Lee and T. Popiel, *The maximal subgroups of the Monster*, submitted (arXiv:2304.14646), 2023.
- [26] J.D. Dixon, *The Fitting subgroup of a linear solvable group*, *J. Austral. Math. Soc.* **7** (1967), 417–424.
- [27] H. Duyan, Z. Halasi and A. Maróti, *A proof of Pyber’s base size conjecture*, *Adv. Math.* **331** (2018), 720–747.
- [28] The GAP Group, *GAP – Groups, Algorithms, and Programming*, Version 4.11.1, 2021 (<http://www.gap-system.org>).
- [29] M. Garonzi and A. Maróti, *On the number of conjugacy classes of a permutation group*, *J. Comb. Theory, Series A* **133** (2015), 251–260.
- [30] R.M. Guralnick and K. Magaard, *On the minimal degree of a primitive permutation group*, *J. Algebra* **207** (1998), 127–145.
- [31] Z. Halasi, *On the base size for the symmetric group acting on subsets*, *Studia Sci. Math. Hungar.* **49** (2012), 492–500.
- [32] Z. Halasi, M.W. Liebeck and A. Maróti, *Base sizes of primitive groups: bounds with explicit constants*, *J. Algebra* **521** (2019), 16–43.
- [33] J.P. James, *Partition actions of symmetric groups and regular bipartite graphs*, *Bull. London Math. Soc.* **38** (2006), 224–232.
- [34] P.B. Kleidman and M.W. Liebeck, *The subgroup structure of the finite classical groups*, *London Math. Soc. Lecture Note Series*, vol 129, Cambridge University Press, 1990.
- [35] R. Lawther, M.W. Liebeck and G.M. Seitz, *Fixed point ratios in actions of finite exceptional groups of Lie type*, *Pacific J. Math.* **205** (2002), 393–464.
- [36] M.W. Liebeck and A. Shalev, *Simple groups, permutation groups, and probability*, *J. Amer. Math. Soc.* **12** (1999), 497–520.
- [37] M.W. Liebeck and A. Shalev, *Bases of primitive permutation groups*, in *Groups, combinatorics & geometry* (Durham, 2001), 147–154, World Sci. Publ., River Edge, NJ, 2003.
- [38] M.W. Liebeck and A. Shalev, *Bases of primitive linear groups II*, *J. Algebra* **403** (2014), 223–228.
- [39] A. Maróti, *Minimal degree, base size, order: selected topics on primitive permutation groups*, *Arch. Math.* **121** (2023), 485–493.
- [40] A. Maróti, *On the orders of primitive groups*, *J. Algebra* **258** (2002), 631–640.
- [41] V.D. Mazurov and E.I. Khukhro, *Unsolved problems in group theory: The Kourovka notebook, no. 20 (English version)* (2022), arxiv:1401.0300.
- [42] G. Mecenero and P. Spiga, *A formula for the base size of the symmetric group in its action on subsets*, *Australas. J. Combin.* **88** (2024), 244–255.
- [43] J. Morris and P. Spiga, *On the base size of the symmetric and the alternating group acting on partitions*, *J. Algebra* **587** (2021), 569–593.
- [44] J. Müller, M. Neunhöffer, F. Noeske and M. Horn, *ORB – methods to enumerate orbits*, Version 4.8.3, 2019. <https://gap-packages.github.io/orb>
- [45] D.S. Passman, *Groups with normal solvable Hall p -subgroups*, *Trans. Amer. Math. Soc.* **123** (1966), 99–111.
- [46] L. Pyber, *Asymptotic results for permutation groups*, in *Groups and Computation* (eds. L. Finkelstein and W. Kantor), DIMACS Series, vol. 11, pp.197–219, 1993.
- [47] Á. Seress, *The minimal base size of primitive solvable permutation groups*, *J. London Math. Soc.* **53** (1996), 243–255.
- [48] E.P. Vdovin, *On the base size of a transitive group with solvable point stabilizer*, *J. Algebra Appl.* **11** (2012), 1250015, 14 pp.
- [49] R.A. Wilson et al., *A World-Wide-Web Atlas of finite group representations*, <http://brauer.maths.qmul.ac.uk/Atlas/v3/>
- [50] R.A. Wilson, *Maximal subgroups of sporadic groups*, in *Finite simple groups: thirty years of the Atlas and beyond*, 57–72, *Contemp. Math.*, vol. 694, Amer. Math. Soc., Providence, RI, 2017.
- [51] R.A. Wilson, *Maximal subgroups of ${}^2E_6(2)$ and its automorphism groups*, preprint (arXiv.1801.08374), 2018.
- [52] R.A. Wilson, R.A. Parker, S. Nickerson, J.N. Bray and T. Breuer, *AtlasRep: A GAP Interface to the Atlas of Group Representations*, Version 2.1.8, 2024. <http://www.math.rwth-aachen.de/~Thomas.Breuer/atlasrep>
- [53] J. Whiston, *Maximal independent generating sets of the symmetric group*, *J. Algebra* **232** (2000), 255–268.

- [54] V.I. Zenkov, *On intersections of nilpotent subgroups in finite symmetric and alternating groups*, Proc. Steklov Inst. Math. **285** (2014), suppl. 1, S203–S208.
- [55] V.I. Zenkov, *On intersections of nilpotent subgroups in finite groups with a sporadic socle*, Algebra Logic **59** (2020), 313–321.
- [56] V.I. Zenkov, *Intersections of three nilpotent subgroups in a finite group*, Sib. Math. J. **62** (2021), 621–637.

M. ANAGNOSTOPOULOU-MERKOURI, SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL, BRISTOL BS8 1UG, UK

Email address: `marina.anagnostopoulou-merkouri@bristol.ac.uk`

T.C. BURNES, SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL, BRISTOL BS8 1UG, UK

Email address: `t.burnes@bristol.ac.uk`